

# Appendix A

## Physics Glossary

This discussion is intended to provide only a passing familiarity with definitions to allow the unfamiliar reader to follow certain calculations and to have a general understanding of the results.

- **absolute space and time:** Newton's view of space and time according to which they exist, independent of whether anything is in the universe or not.

- **abstract spin network:** They are the equivalence classes of spin networks under spatial diffeomorphisms. GR tells us that the spacetime manifold is a convenient mathematical device devoid of physical meaning. How a spin network is embedded in the spacial three-slice. Once we factorize out the spatial diffeomorphism invariance the only remaining information in the state is the knottedness of the loop. These are objects that don't live in the manifold anymore - they are not embedded in a space. They are not quantum excitations in space, but quantum excitations of space - if there is no spinor network there is no space!

- **achronal:** A subset  $S$  of spacetime  $\mathcal{M}$  is said to be achronal provided no two of its points can be joined by a timelike curve.

- **active diffeomorphism:** transformation relates different objects in  $\mathcal{M}$  in the same coordinate system. This means that the diffeomorphism  $f$  is viewed as a map associating one point of the manifold to another one.

Invariants with respect to active diffeomorphisms are obtained by solving away the coordinates  $x$  from solutions to the equations of motion.

is a solution of Einstein's equations, namely active diffeomorphisms are dynamical symmetries of Einstein's equations sending solutions to solutions.

- **ADM:** In the Hamiltonian formulation of GR the generator of time translations in an asymptotically flat spacetime.

(1) ADM energy

(2) ADM angular momentum

- **Alexandroff topology:** For a spacetime  $\mathcal{M}$  and any two spacetime points  $a, b$ . The chronological sets  $\{I^+(a) \cap I^-(b) : a, b \in \mathcal{M}\}$  are open and form a base for a topology on  $\mathcal{M}$ , which is called the Alexandroff topology. (Also see maths glossary).

- **algebraic quantum theory:** In the Heisenbergs picture of quantum mechanics the observables play the dominant role, they are governed by an equation of motion while the states are time-independent. In algebraic quantum theory we formulate more abstract version of Heisenberg's approach, in which the observables are the relevent objects and the quantum states are secondary.

There are two main ingredients: firstly, a  $C^*$ -algebra of observables often denoted  $\mathcal{A}$ , and secondly, states (different meaning from above)  $\omega$  which act on an observable  $A \in \mathcal{A}$  giving a complex number i.e.  $\omega : \mathcal{A} \rightarrow C$  (In this report we use the notation  $E_\psi$  instead of  $\omega$ ),

$$E_\psi(\cdot) := \frac{\langle \psi, \cdot \psi \rangle}{\|\psi\|} \iff \omega(\cdot) \quad (\text{A.1})$$

These states are positive linear functionals ( $\omega(A^*, A) \geq 0$  for all  $A \in \mathcal{A}$ ) such that  $\omega(I) = 1$ , or in our notation

$$E_\psi(A) = \frac{\langle \psi, A^* \hat{A} \psi \rangle}{\|\psi\|} \geq 0 \text{ for all } A \in \mathcal{A}, \quad E_\psi(I) = \frac{\langle \psi, \hat{I} \psi \rangle}{\|\psi\|} \equiv 1. \quad (\text{A.2})$$

The value of the state  $\omega$  acting on an observable  $A$  can be interpreted as the expectation value of the operator  $A$  on the state  $\omega$ , i.e.  $\langle A \rangle = \omega(A)$ , or

$$E_\psi(A) = \frac{\langle \psi, \hat{A} \psi \rangle}{\|\psi\|} \quad (\text{A.3})$$

- **affine quantum gravity:** [402]

- **algebraic quantum field theory (AQFT):** Reformulating QFT on an axiomatic basis: that is, starting from what seem to be physically necessary and mathematically precise principles which any QFT would have to satisfy, and then finding QFTs which actually satisfy them, (see [114], and references therein, for extensive discussion of it).

Fields at a point. In the formulation of quantum field theory one traditionally starts with a particle picture and investigate the scattering theory.

takes the collection of all operators localized in a particular region.

Operators of observables localized in an open region of spacetime  $\mathcal{O}$  form an algebra  $\mathcal{A}(\mathcal{O})$  of bounded operators. It is possible to encode all physically relevant properties in terms of these algebras and their transformation behaviour under Poincaré group.

Algebraic quantum field theory (see [1]) is a general approach to quantum field theory based on algebras of local observables, the relations among them, and their representations.

Haag-Kastler axioms

- (i) powerful structural analysis proven useful for rigorous treatment of models,
- (ii) revision of the particle concept,
- (iii) algebraic approach has shed new light on the treatment of quantum field theories on curved spacetime.

the physical concept of locality of nets of local algebras  $\mathcal{O} \mapsto \mathcal{A}(\mathcal{O})$ . For each open region  $\mathcal{O}$  one assigns a  $C^*$ -algebra  $\mathcal{A}(\mathcal{O})$ . These are mutually (anti)commuting for spacelike separated regions.

In AQFT one cleanly separates two states of quantizing a field theory, namely first to define a suitable algebra  $\mathcal{A}$  and then study its representations in a second step.

The axiomatic nature of AQFT makes it ideal for foundational study, since we can specify with mathematical precision exactly what the entity is that we are studying.

• **algebraic quantum gravity:** Algebraic quantum gravity, proposed by Thiemann [2], is a new approach canonical quantum gravity suggested by loop quantum gravity.

It has been shown that algebraic quantum gravity admits a semiclassical limit whose infinitesimal gauge symmetry agrees with that of general relativity.

• **algebra of kinematic observables:** kinematic observables because they are not gauge invariant. These separate points of the unconstrained classical phase space of the system - that is  $\mathcal{P}$ . More complicated composite functions can be expressed in terms of (of limits) them. Dirac constraint approach: We want to ultimately construct a physical Hilbert space of solutions to the constraints. The kinematic observables can be projected onto the physical observables.

In the reduce phase space scheme we first find the physical Dirac observables and then attempt to quantize our system by finding a representation of the Poisson algebra of these observables as linear operators on a Hilbert space.

• **almost periodic functions:**

• **alternative theories of quantum gravity:**

Well developed background independent approaches to quantum gravity are causal sets, causal dynamical triangulations and quantum Regge calculus.

- **anomalies:** When we quantize a classical system it is possible that a symmetry of the classical theory is lost. That is, an anomaly is a classical symmetry not carried over to the quantum theory. Anomalies arise when the classical brackets cannot be fully realized by the quantum commutators.

- **apparent horizon:** See trapped surface.

i)  $S$ , the expansion  $\Theta_{(\ell)}$  of the one null normal,  $\ell^a$ , is everywhere zero and that

ii) that of the other null normal,  $n^a$ , is negative.

- **Arnowitt-Deser-Misner (ADM):** together, (as well as Dirac), they laid much of the groundwork for canonical quantum gravity. They expressed GR in Hamiltonian form by splitting spacetime into  $R \times \sigma$  where  $R$  (the real line) represents “time” and  $\sigma$  is a three-dimensional space at a given “time”. As their “configuration” variable they used the 3-metric that the spatial slice picks up from the metric of the spacetime, and as their “conjugate momentum” the extrinsic curvature(-to do with the time derivative of the 3-metric). This Hamiltonian construction is known as the ADM formalism.

- **Ashtekar, Abhay:**




Figure A.1: Abhay Ashtekar.

- **Ashtekar’s new variables:** A major obstacle to progress in the canonical approach had been the complicated nature of the field equations in the traditional variables,  $(q_{ab}, p^{ab})$ . This obstacle was removed in 1984 with the introduction of new canonical variables - Ashtekar’s new variables. In terms of these, *all* equations of the theory become polynomial, in fact, at worst quartic, [93].

Self-dual connection. This odd choice leads to constraint equations that are polynomial in the basic variables - the connection  $A$  and the “electric” field  $E$ . The Hamiltonian constraint, that was non-polynomial in the ADM treatment, is quadratic in  $A$  and  $E$ .

- **Ashtekar-Isham-Lewendowski state:**

Haar measure on cylindrical functions.

- **asymtotically flat gravitational field:**

an asymptotically flat gravitational field: in this case the Hamiltonian is given by suitable boundary terms and the observables at infinity evolve in the Lorentz time of the asymptotic metric.

- **asymptotic flatness:** There exists coordinates  $x = (x_1, x_2, x_3)$  defined outside a compact set on  $\Sigma$  such that, for some  $\delta > 1$ ,

$$h_{ij} - (1 + \frac{2M}{r})\delta_{ij} = \mathcal{O}(r^{-\delta}),$$

as  $r \rightarrow \infty$  with  $M \geq 0$  (positive mass theorem).

- **automorphisms:** generalized Heisenberg picture one-parameter group of automorphisms of the algebra of the algebra of observables. Take any  $A \in \mathcal{A}$  an automorphism is

$$\gamma_t A = e^{it\hat{\mathcal{H}}/\hbar} A e^{-it\hat{\mathcal{H}}/\hbar} \tag{A.4}$$

where  $\gamma_t A \in \mathcal{A}$ . Obviously

$$\gamma_s \gamma_t A = \gamma_{s+t} A \tag{A.5}$$

$$\frac{d}{dt} A(t) = [A(t), \mathcal{H}]. \tag{A.6}$$

- **background independent:** There are various ways of expressing this property - it means that how a particle or a field is localization with respect to a spacetime manifold has no physical significance.

The theory does not depend on a background metric, thus all distances are gauge equivalent.

There is no fixed geometric component to the Einstein-Hilbert action.

- **back-reaction:** in some background spacetime affect the dynamics of the background. This back-reaction can be described in terms of an effective energy-momentum tensor  $\tau_{ab}$ .

$$G_{\mu\nu} = \kappa \langle \hat{T}_{\mu\nu} \rangle \quad (\text{A.7})$$

Gravitat

Dynamical theory over curved spacetime not a dynamical theory of spacetime.

see quantum field theory on curved spacetime.

- **Baez, John**




Figure A.2: John Baez.

- **Barbour, Julian:** Reinterpreted Einstein's general theory of relativity as a relational theory in which space and time are nothing but a system of relations. Since receiving his doctorate in 1968 from the University of Cologne, has never held an academic job but became an honored member of the quantum gravity community as it was Barbour who taught people what it means to make a background-independent theory.

- **Bargmann representation:**

- **Barrett, John**

- **Barrett-Crane model:** The Barrett-Crane model is one of the most extensively studied spin foam models for quantum gravity. It can be viewed as a spin foam quantization of the  $SO(4)$  Plebanski formulation of general relativity. Specifically, one quantizes the Topological BF theory then imposes the simplicity constraints (that convert BF theory into GR) at the quantum level.



Figure A.3: Julian Barbour.

The original strategy followed by Barrett and Crane the starting point was with the quantization of a 4-simplex.

There are problems with the Barret-Crane models from its ultra-locality which follows from the way in which the simplicity constraints are imposed. Various proposals have been made recently to address these problems, [200], [286] . From [286]: “... that a new exciting period of the development of the subject of spin foam models may be opening.”

- **baryonic matter:** Baryonic matter or simply ordinary matter is anything made of atoms and their constituents, and this includes all of stars, planets, gas and dust in the universe. Ordinary baryonic matter, it turns out, is not enough to account for the observed matter density; Baryons are present in the amount predicted by the Big Bang Nucleosynthesis, some percent of the density required to close the universe.

- **Benny Hill effect:** There is a negative force between a particle of negative mass and a particle of positive mass, whereas by  $F_n = m_n a_n$  ( $F_n$  is the force exerted on the negative mass particle and  $a_n$  is the acceleration of the negative particle mass particle) the negative mass particle accelerates towards the positive mass particle, while the positive particle mass accelerates away.

- **Bergmann-Komar group:** Transformations between different covariant gauge fixings form a group - Bergmann-Komar group.

- **beta function:**

- **Bekenstein bound:**



Figure A.4: John Barrett.

$$S < 8\pi\zeta RE \tag{A.8}$$

- **Big bang:**

journal Advances in Theoretical and Mathematical Physics:

*“The question of whether the universe had a beginning at a finite time is now ‘transcended’. At first, the answer seems to be ‘no’ in the sense that the quantum evolution does not stop at the big-bang. However, since space-time geometry ‘dissolves’ near the big-bang, there is no longer a notion of time, or of ‘before’ or ‘after’ in the familiar sense. Therefore strictly, the question is no longer meaningful. The paradigm has changed and meaningful questions must now be phrased differently, without using notions tied to classical space-times.”*

- **Biot-Savart law:**

$$B = -\frac{1}{4\pi} \int d^3y \epsilon_{\alpha\beta\gamma} j^\beta(y) \frac{(x-y)^\gamma}{|x-y|^3} \tag{A.9}$$

- **black-body radiation:** Thermal radiation emitted by black-bodies, black-bodies being perfect emitters and absorbers of radiation. photons moving around randomly, without any discernible source.

- **black hole entropy:**

- **black hole:** collapsing star forming an object so dense that nothing can escape from it. A region of spacetime hidden behind a horizon.




Figure A.5: Lebrun.

- **black hole uniqueness theorems:** If a black hole is stationary, then it must be either static or axially symmetric. This is the Kerr family of black holes
- **Bochner-Minlos theorem:** Fourier transform corresponds a unique positive-definite measure.
- **Bogoliubov transforms:** Two different Fock space representations are related by a transformation to new operators.

In ordinary quantum field theory in flat spacetime, one quantizes  $\phi$  by first decomposing the field into Fourier modes,

$$\phi = \sum_{\mathbf{k}} \left( a_{\mathbf{k}} u_{\mathbf{k}}(t, \mathbf{x}) + a_{\mathbf{k}}^{\dagger} u_{\mathbf{k}}^*(t, \mathbf{x}) \right)$$

with

$$u_{\mathbf{k}} = e^{i\mathbf{k}\cdot\mathbf{x} - i\omega_{\mathbf{k}}t}, \quad \omega_{\mathbf{k}} = (|\mathbf{k}|^2 + m^2)^{1/2}$$

and replacing the coefficients  $a_{\mathbf{k}}, a_{\mathbf{k}}^{\dagger}$  by annihilation and creation operators respectively. The Fourier modes  $u_{\mathbf{k}}$  are a set of orthonormal functions satisfying

$$(\square + m^2)u_{\mathbf{k}}(t, \mathbf{x}) = 0, \quad \partial_t u_{\mathbf{k}}(t, \mathbf{x}) = -i\omega_{\mathbf{k}} u_{\mathbf{k}}(t, \mathbf{x}),$$

where the second condition determines what one means by positive and negative frequency, and thus allows to distinguish creation and annihilation operators.

In a curved spacetime, or noninertial reference frame in flat spacetime, standard Fourier modes are no longer available. However, with a choice of time coordinate  $t$  one can still find a decomposition to obtain creation and annihilation operators. Given two different reference frames with time coordinates  $t$  and  $\bar{t}$ , two such decompositions exist:

$$\phi = \sum_i (a_i u_i + a_i^\dagger u_i^*) = \sum_i (\bar{a}_i \bar{u}_i + \bar{a}_i^\dagger \bar{u}_i^*)$$

and since  $(u_i, u_i^*)$  are a complete set of functions, we can write

$$\bar{u}_j = \sum_i (\alpha_{ji} u_i + \beta_{ji} u_i^*).$$

This relation is known as a Bogoliubov transformation, and the coefficients  $\alpha_{ji}$  and  $\beta_{ji}$  are called Bogoliubov coefficients.

- **Bojowald, Martin:**




Figure A.6: Martin Bojowald.

- **Boltzmann statistical mechanics:**

- **Bondi coordinates:**  $\mathcal{I}^+$  can be foliated into a one parameter family of two dimensional non-intersecting spacelike surfaces, each diffeomorphic to  $S^2$ , parametrized by a variable

$u$ . On each of these surfaces one can introduce complex stereographical coordinates  $\zeta$  and  $\bar{\zeta}$ . The coordinates  $(u, \zeta, \bar{\zeta})$  on  $\mathcal{I}^+$  are called the Bondi coordinates on  $\mathcal{I}^+$ .




Figure A.7: Bondi coordinates at future null infinity.

- **Bondi-Sachs mass:** Both the energy of the radiation and the energy of the leftover system are included in the total ADM energy. The Bondi energy is the gravitating mass as seen by light rays propagating out to infinity in the lightlike direction, rather than the spacelike direction. For a spacetime of an isolated body emitting gravitational the Bondi-Sachs mass decreases, while the ADM mass stays constant.

- **Born-Oppenheimer approximation:**

- **bottom-up:** ‘bottom-up’ approach one attempts to formulate the full theory and then make some approximation to get the low energy description. Hopeful, in doing so we recover the correct classical theory. When there are an infinite number of degrees of freedom, the passage from the quantum theory to the classical regime does not always have to give the classical theory that was quantized!

- **BRST:** The central idea of BRST theory is to replace the original gauge symmetry by a rigid fermionic symmetry acting on an appropriately extended space containing new variables, “ghosts”. Ghosts first appeared in quantum field theory - they ensured that the theory would be unitary and independent of the gauge choice. They did this through contributions to virtual process only - they have the wrong spin statistics to correspond to physical particles. They violate the spin statistic relation which requires physical fields with half-integer spin to be fermionic and fields with integer spin to be bosonic.

- **bubble formulism:** Kuchař’s bubble time formulism

- **canonical:** Simplest or standard form.

- **canonical quantisation:** Classical mechanics admits two different kinds of formulation: Hamiltonian and Lagrangian. So does quantum mechanics. The one corresponding to the Hamiltonian formulation of classical mechanics is called canonical quantisation and involves a Hilbert space and operators.

The basic idea is

- (i) to take the states of the system to be described by wavefunctions  $\Psi(q)$  of the configuration variables,
- (ii) to replace each momentum variable  $p$  by differentiation with respect to the conjugate configuration variable, and
- (iii) to determine the time evolution of  $\Psi$  via the Schrödinger equation,

$$i\hbar \frac{\partial \Psi}{\partial t} = \hat{\mathcal{H}}\Psi,$$

where  $\hat{\mathcal{H}}$  is an operator corresponding to the classical Hamiltonian  $\mathcal{H}(p, q)$ .

General relativity can be cast in Hamiltonian form, however the Hamiltonian is a constraint on phase space. Dirac developed the general method for the canonical quantisation of systems with constraints. It involves the imposition of the constraints as an additional condition on the Hilbert space.

- **canonical transformations:** A canonical transformation in a classical theory relates descriptions by different ‘generalised coordinates’ on phase space with the same symplectic structure. These classical theories are equivalent, but the resulting quantum theories need not be equivalent.
- **Cartan’s structure equations:** See maths glossary.
- **Cauchy Horizon:**  $D^-(p)$




Figure A.8: Cauchy Horizon.

- **Cauchy surface:** A Cauchy surface is a spacelike or null surface that intersects every timelike curve in a spacetime  $\mathcal{M}$  once and only once. (Not every spacetime has a Cauchy surface! - see global hyperbolicity).
- **causal** As the very concepts of time and spacial location are not fundamental at all in quantum gravity and only emerges in a semiclassical approximation, causality should too only emerge and be applicable only in a semi-classical limit.

??but no background causal structure in the classical theory - the notion of causality only emerges when the dynamics of gravity can be neglected??

- **causal future:** events are influenced by those in their past.
- **causality conditions:** Avoid paradoxes traveling into the past. Heierarky of conditions:

- 1) Chronology: no closed timelike curves;
- 2) Causality: no closed causal curves;
- 3) Strong causality: no “almost closed” causal curves;
- 4) Stably causality: close metrics are causal.

- **causal set theory:** The causal approach to quantum gravity is the proposal that the structure of spacetime at the quantum scale should be modelled after a locally finite partially ordered set (poset) of elements, the *causal set*, whose partial order is the relation that defines past distinction between events.

See quantum causal histories.

[gr-qc/0507078]. This approach does not attempt a quantization of classical theory. Rather, it is a construction of a quantum theory of causal sets based on two features of classical genral relativity that it taakes as fundamental: the causal structure, which is replaced by discrete causal set;

- **causal spin networks:**

Do not necessarily need to build in causality as it could be an emergent property??

- **CBR, cosmic background radiation:** Thermal radiation. photons moving around randomly, without any discernible source, corresponding to a temperature of about  $2.7K$ .

- **charge spin networks:**  $U(1)$  gauge group labelled by integers.

- **Chern-Simons:**

Relevant in condensed-matter physics - description of excitations with fractional statistics, *anyons*, in the fractional Hall effect (FQHE) and a proposal for a mechanism of superconductivity based on anyons Y-H. Chen, F. Wilczek, E. Witten, and B.I. Halerin, Int. J. Mod. Phys. B **3**, 1001 (1989).

- **Chern-Simons-Witten gravity:** The local Poincare  $ISO(2, 1)$  symmetry is, on shell, equivalent to the usual diffeomorphism [387]. This  $(2 + 1)$  dimensional gravity is based on the non-degenerate, invariant and associative inner product on  $iso(s, 1)$  on which the conventional Killing metric is degenerate [387]:

How do we define the trace operation in Chern-Simons define

$$\langle P^a, J^a \rangle = \eta^{ab}, \quad \langle J^a, J^b \rangle = \langle P^a, J^b \rangle = 0, \quad (\text{A.10})$$

$$[P^a, J^a] = \epsilon_{abc} J^c, \quad [J^a, J^b] = \epsilon_{abc} P^c \quad [P^a, P^b] = 0, \quad (\text{A.11})$$

where  $\eta^{ab} = \text{diag}(-1, 1, 1)$  and  $P^a, J^a$  are the generators of  $iso(2, 1)$  (here Roman indices are for internal coordinates and Greek indices for spacetime coordinates). The gauge connection is an  $iso(2, 1)$  valued one form

$$A = \omega_a J^a + e_a P^a, \quad (\text{A.12})$$

where  $\omega$  and  $e$  are the homogenous (Lorentz) and the inhomogeneous (translational) part of the connection components respectively.

ISO(2,1) Gauge theory with the gravitational Chern-Simons term

If we define  $\text{Tr}(\cdot) := \langle \cdot, \cdot \rangle$  so that (A.11)

$$\text{Tr}(P^a J^a) = \eta^{ab}, \quad \text{Tr}(J^a J^b) = \text{Tr}(P^a J^b) = 0, \quad (\text{A.13})$$

$$\int \text{Tr} \left\{ A \wedge dA + \frac{3}{2} A \wedge A \wedge A \right\} \quad (\text{A.14})$$

- **chromodynamics:**

- **closed systems:** The universe as a whole. while the standard Copenhagen interpretation works well in the laboratory, it cannot be applied to closed systems.

- **classical configuration space:** The classical configuration is generally taken to be the space of all smooth functions which decay rapidly at infinity on a  $t = \text{constant}$  slice.

- **clock variable:** A clock variable is a function on the extended phase space of a system which strictly monotonous along the evolution orbits...

- **closed trapped surface:** See trapped surfaces.

- **coherent states:** They provided the closest approximation to classical physics as uncertainty is minimum and the states are peaked in both configuration and momentum representations.

- **collapse of the wave function:**

- **Coleman-Mandula theorem:** The theorem states that a group that non-trivially combines both the Lorentz group and a compact Lie group cannot have finite dimensional,

unitary representations. This presents a problem to constructing unified theory of gravity and the other forces. A way to evade the Coleman-Mandula theorem is to use super symmetry.

- **comparative property:** a comparative property and to express it we need a way to introduce scale into our system.

- **constraints:**

we can take these constraints to say that small automorphisms of the bundle  $P|_S$

- **constructive quantum field theory:**

Geared to scalar fields propagating on Minkowski space. Key axioms are called the Osterwalder-Schrader axioms.

- **common eigenstate:**

- **complete set of commuting operators:** In three-dimensions, three coordinate operators  $\hat{x}, \hat{y}$  and  $\hat{z}$ . As they commute with each other we can measure one then measure the next without changing the . and uniquely specifies the state of the system.

The outcomes of the measurement of a complete set of commuting observables characterizes the state.

When we have spherical symmetry potentials in three-dimensions, the set  $\hat{\mathcal{H}}, \hat{L}^2$  and  $\hat{L}_z$

We can define an orthonormal basis in this Hilbert space by diagonalizing a complete set of commuting self-adjoint operators.

Measuring  $A$  first, we would find a value  $a_n$  and would leave the system in an eigenstate of  $A$ .

$$A|\psi_i \rangle = a_i|\psi_i \rangle, \quad B|\psi_i \rangle = b_i|\psi_i \rangle \quad (\text{A.15})$$

$$(AB - BA)|\psi_i \rangle = 0 \quad \implies \quad AB|\psi_i \rangle = a_i B|\psi_i \rangle \quad (\text{A.16})$$

$$\langle \psi_i | (AB - BA) | \psi_j \rangle = (a_i - a_j) \langle \psi_i | B | \psi_j \rangle = 0, \quad (\text{A.17})$$

So  $\langle \psi_i | B | \psi_j \rangle$  vanishes unless  $a_i = a_j$ .

$|\psi_i \rangle$  are **common** eigenstates of  $\hat{A}$  and  $\hat{B}$ . We denote them as  $| \rangle$

Measuring  $A$  first we find it has the value  $a_i$  and the system would be left in the eigenstate of  $A$ . As shown above, this eigenstate is also an eigenstate of  $B$ . Thus a measurement of  $B$  yields a value  $b_i$

**Definition:** A set of self-adjoint operators is called a CSCO if the operators all commute with each other and if the set of their common eigenstates is complete and not degenerate.

- **complexifier:** The “complexifier” was introduced by Thiemann in [243] as a powerful tool that can give rise to kinematical coherent states for a general quantum system.

the complexification of  $SU(2)$  is  $SL(2, \mathbf{C})$ .

$$u = \exp[i\tau_j^j/2] \tag{A.18}$$

- **Compton scattering:** When a photon scatters off a free electron, in the relativistic case  $\hbar f \gg m_e c^2$ .

- **conformally flat spacetimes:**

$$g_{ab} = \Omega \eta_{ab} \tag{A.19}$$

- **conformal transformations:**

- **congruence:** A congruence is a family of curves such that precisely one curve of the family passes through each point. It is a geodesic congruence if the curves are geodesics. A vector field  $v^a$  determines a congruence whose curves are tangent to  $v^a$ . We call such curves streamlines of  $v^a$ .

- **conjugate momentum:**

$$p_i := \frac{\partial \mathcal{L}}{\partial q^i}. \tag{A.20}$$

$$\{q, p\} = 1 \tag{A.21}$$

In the Hamilton formulation of GR, the conjugate momenta of the induced metric is related to the extrinsic curvature  $K^{ab}$  by

$$\pi^{ab} = q^{-1/2}(K^{ab} - Kq^{ab}). \tag{A.22}$$

- **connection:** self-consistent way of comparing vectors in two different vector spaces.

We define connection and curvature variables  $A_{(\gamma)\mu}^{IJ}$  and  $F_{(\gamma)\mu\nu}^{IJ}$

$$A_{\mu}^{IJ} = \omega_{\mu}^{IJ} + \frac{\gamma}{2} \epsilon^{IJ}{}_{KL} \omega_{\mu}^{IJ} \tag{A.23}$$

$$F_{\mu\nu}^{IJ} = R_{\mu\nu}^{IJ} + \frac{\gamma}{2} \epsilon^{IJ}{}_{KL} R_{\mu\nu}^{KL} \quad (\text{A.24})$$

If one chooses the value of  $\gamma$  such that  $\gamma^2 \sigma = 1$ , then the variables  $A_{\mu}^{IJ}$  and  $F_{\mu\nu}^{IJ}$  are self-dual with respect to the internal indices.

- **Connes-Kreimer's Hopf algebra of Feynman's diagrams:** renormalization via Hopf algebra of rooted trees. Applied by Fontini Markopoulou to renormalization of spin foams background-independent models of quantum gravity.

- **consistent discretizations:** Rodolfo Gambini Jorge Pullin based on approximating the continuum theory by a discrete theory. It allows to bypass difficult conceptual problems turning them into technical ones.

relational quantum theories unfortunately one would have to disregard all the results of LQG

**consistent discrete quantum gravity *a la* Pullin, Gambini *et al*:** A discrete approach to quantum gravity that is free of constraints translating difficult conceptual problems into problems of computational nature.

- **consistent-histories approach to quantum theory** The consistent-histories approach to quantum theory was originally introduced to provide a novel way of re-interpreting standard quantum theory, particularly in regard to the role played by measurement. However, because of the novel way in which time is handled, consistent-history theory also has the potential for providing new and powerful ways of studying quantum theories of gravity. Most recently, in [2] the formalism was applied to construct a history version of the canonical form of classical general relativity. The possibility also arises to use this formalism in the context of generalised ideas of time and space: for example, in models where spacetime is not represented by a differentiable manifold.

[?] 28 Oct 2001, *Quantising the Foliation in History Quantum Field Theory* C.J. Isham, K. Savvidou, [quant-ph/0110161].

- **constants of motion:**

$$\frac{d}{dt} \langle \hat{A} \rangle = \frac{1}{i\hbar} \langle [\hat{\mathcal{H}}, \hat{A}] \rangle + \langle \frac{\partial \hat{A}}{\partial t} \rangle . \quad (\text{A.25})$$

If  $\hat{A}$  commutes with the Hamiltonian and does not explicitly depend on time, the observable  $A$  is said to be a constant of motion.

$$[\hat{\mathcal{H}}, \hat{A}] = 0 \quad (\text{A.26})$$

- **constraint:** [89]: In addition to the Hamiltonian equations of motion, the theory will exhibit constraint equations. The constraints play a double role in a Hamiltonian theory.

On the one hand they generate a group of symmetries of the phase space referred to as the gauge transformations, on the other hand the set of solutions of the constraints defines the constraint surface of the phase space.

The simplest example for such a kind of theory is certainly Maxwell theory, where the structure group is  $U(1)$ . A more general example is Yang Mills theory, where the structure group may be an arbitrary compact Lie group  $G$ . In this case the group of the gauge transformations is the group of the fiber preserving automorphisms of the given bundle, homotopic to the identity. The group is often referred to as the Yang-Mills gauge transformations.

• **constraint algebra:**

quantum constraint algebra - is mathematically consistent but seems to fail to reproduce the classical constraint algebra - however in calculating the righthand side of the classical Hamilton-Hamilton Poisson bracket involves... So not obvious if it does fail??

• **constructive quantum field theory:** The program of constructive quantum field theory was initiated in the early seventies. This approach has had remarkable success in certain 2 and 3 dimensional models. From a theoretical physics perspective, the underlying ideas may be summarized roughly as follows. Consider, for definiteness, a scalar field theory. The key step then is that of giving meaning to the Euclidean functional integrals by defining a rigorous version  $d\mu$  of the heuristic measure “[ $\exp(S(\phi))$ ]  $\prod_x d\phi(x)$ ” on the space of histories of the scalar field, where  $S(\phi)$  denotes the action governing the dynamics of the model. The appropriate space of histories turns out to be the space  $\mathcal{S}'$  of (tempered) distributions on the Euclidean space-time and regular measures  $d\mu$  on this space are in one to one correspondence with the so called generating functionals, which are functionals on the Schwarz space  $\mathcal{S}$  of test functions satisfying certain rather simple conditions. (Recall that the tempered distributions are continuous linear maps from the Schwarz space to complex numbers.)

Thus, the problem of defining a quantum scalar field theory can be reduced to that of finding suitable measures  $d\mu$  on  $\mathcal{S}'$ , or equivalently, appropriate functionals  $\chi$  on  $\mathcal{S}$ . Furthermore, there is a succinct set of axioms due to Osterwalder and Schrader which spells out the conditions that  $\chi$  must satisfy for it to be appropriate, i.e., for it to lead to a consistent quantum field theory which has an associated

- i. Hilbert space of states,
- ii. a Hamiltonian,
- iii. a vacuum and
- iv. an algebra of observables.

(from Integration on the space of Connections Modulo Gauge Transformations)

For introduction see for example [397].

• **coordinate singularity** Are locations in coordinate space where calculations become degenerate or divergent, due solely to “bad coordinates”. An example of this is the

Schwarzschild radius  $r = 2M$  in Schwarzschild coordinates. A radially-infalling observer takes an infinite amount of time in the Schwarzschild time coordinate to cross the Schwarzschild radius. The proper time of the observer would measure although he would see the outside universe speed up?????

One can avoid the detection of unreal coordinate singularities by maximally extending the space-time.

- **Copenhagen interpretation:** ??? (quant-ph/9803052) an a priori classical part of the world is assumed to exist from the outset. Such a structure is there thought to be necessary for the "coming into being" of observed measurement results. ???

In the absence of a background spacetime geometry, the Copenhagen interpretation is of little use - it is definitely physically incomplete.

[338] "The usual well defined quantum operators like the area and volume will exist and be well defined, except that in the discrete theory the total volume of the slice will be an observable since there are no further constraints."

no external observer of the universe as a whole throws into doubt the instrumentalism of the Copenhagen interpretation; as does an attempt to construct a quantum gravity theory with no background spacetime in which an 'observer' could be placed. [362].

standard Copenhagen divide the physical world into the microscopic regime that obeys the laws of quantum mechanics and the macroscopic regime which obeys the laws of classical mechanics e.g. Maxwell's equations, Einstein's equations. Observers are classical. artificial partition not clear where to put this quantum/classical border. (quant-ph/0103161)

- **Corichi, A:**

- **cosmic censorship:** The principle that singularities are never "naked", that is, do not occur unless surrounded by a shielding event horizon.

- **cosmological constant:** A constant introduced into Einstein's field equations of general relativity in order to supplement to gravity. If positive (repulsive), it counteracts gravity, while if negative (attractive), it augments gravity. It can be interpreted physically as an energy density associated with space itself.

- **cosmological constant problem:** zero point energy of quantum fields - why is  $\Lambda$  not infinite? With a Plankian cut-off the value obtained from calculations performed in quantum field theory on the vacuum energy density corresponding to quantum fluctuations of the fields we observe in nature, imply values that are over 120 orders of magnitude greater than the observed value.

In his model, he was able to show he cosmological constant paradox appears only if spacetime is regarded as fundamental rather than emergent, [266]. The excitations are part of the pre-geometry and cannot curve the effective spacetime they simulate. The

fields we observe in nature emerge as in the low energy, coarse grained limit of a more fundamental theory.

- **cosmological principle:** this states that the universe must appear the same whatever the point from which it is observed. It implies that the universe is homogeneous and isotropic on the largest scales and forms the foundations of all of present day cosmology.

- **course graining:** A nice example of a Hopf algebra.

background-independent course graining *a la* Fontini-Markopoulou [122], [?]

- **covariant quantum statistical mechanics:**

intensive thermodynamical quantities extensive thermodynamical quantities

- **Covariant Loop Quantum Gravity:** (CLQG) is a loop quantization of a Lorentz covariant formulation. It uses the full Lorentz gauge group and produces results independent of the Immirzi parameter.

- **Crane, L:**

- **crossing-symmetry:** In gravity this has a relation to the following: [151], [23]. 4d-diff invariant spinfoam. A problem of the canonical formalism is that it is necessary to make a  $(3 + 1)$  decomposition of spacetime which breaks the manifest 4-diffeomorphism invariance. The vector constraint, generated space diffeomorphisms, and the scalar constraint, generating time reparametrization, arise from the underlying 4-diffeomorphism invariance. It reduces the full, 4-dimensional invariance - the canonical formalism preserves spatial diffeomorphism invariance within each hypersurface of the foliation.

There is a suggestion for curing the slicing dependence.

the spacetime covariant formalism in [151] naturally suggest a “covariantisation” of the operator, described in [151] under the name of “crossing symmetry”.




Figure A.9: crossing. The crossing symmetry vertices.

Actually, a well known property of standard QFT is that the vertex amplitude does not depend on which states are “in” and which are considered “out” - this is related to the crossing symmetry mentioned here.

- **curvature:**

- **curvature tensor:** The vector  $X$  defines a curve through the point  $p$  via parallel transport, The vector  $Y$  defines another curve through  $p$ . We can form an attempted parallelogram.




Figure A.10: We display the geometric interpretation of the curvature tensor. Carry a third vector  $Z$ , by parallel transport from  $p$  to  $s$  via  $q$ , comparing this with transporting this from  $p$  to  $s'$  via  $r$  we find a discrepancy between the two vectors given in terms of the curvature tensor components  $R_{abc}{}^d$  by the formula  $\epsilon^2 X^a Y^b Z^c R_{abc}{}^d$ .

- **cut-off coherent states:** The coherent state  $\psi_m$ , produced by the complexifier method, with respect to a finite graph as a graph dependent coherent state in  $\mathcal{H}_{Kin}$  is called a cut-off coherent state. These are then normalizable, graph dependent states, however they can not be used to test the semiclassical properties of graph changing operators, such as the Hamiltonian constraint, as their expectation values with respect to these cut-off states is always zero.

- **dark matter:** A kind of particle that may exist in our universe. This may not be part of the standard Model of particle physics.

conclusion comes from galaxy rotation curves,

[?] A proper General relativist calculation shows a large departure from Newtonian approximation,

*A galaxy is modelled as a stationary axially symmetric pressure free-fluid in general relativity. For weak gravitational fields under consideration, the field equations and the equations of motion ultimately lead to one linear and one non-linear equation relating the angular velocity to the fluid density. It is shown that the rotation curves for the Milky Way, NGC 3031, NGC 3198 and NGC 7331 are consistent with the mass density distribution of the visible matter cocentred in flattened disks. Thus the need for massive*

*halo of exotic dark matter is removed. For these galaxies we determine the mass density for the luminous threshold as  $10^{-21.75} \text{ kg}\cdot\text{m}^{-3}$ .*

“It is understandable that the convectional gravity approach has focused upon Newtonian theory in the study of galactic dynamics as the galactic field is weak (apart from the deep core regions where black holes are said to reside) and the motions are non-relativistic ( $v \ll c$ ).”

“We have seen that the non-linearity for the computation of density inherent in the Einstein field equations for a stationary axially-symmetric pressure-free mass distribution, even in the case of weak fields, leads to the correct galactic velocity curves as opposed to the incorrect curves that had been derived on the basis of Newtonian gravitational theory.”

• **decoherence:** what decoherence does for you is make any discrepancies minute so we don't have to worry about conceptual difficulties. alleviates a practical problem - it is not clear if it elevates conceptual difficulties.

kent dowker (1995): one might at least hope to predict the gross features: the existence and persistence of (what appears to us to be) a classical world with very large-scale structure; the fact that matter clumps and fields in this classical world generally obey classical equations of motion to a very good degree of approximation;

interacting with the classical world, follow the probabilistic laws of Copenhagen quantum theory.

• **decoherent histories:** Taken from - Quantum logic and decohering histories, [quant-ph/9506028]:

In recent years, much attention has been devoted to the so-called decoherent histories approach to quantum theory. A major motivation for this scheme is a desire to replace the traditional Copenhagen interpretation of quantum theory with one that avoids any fundamental split between observer and system and the associated concept of state-vector reduction induced by a measurement. The key ingredient of the new approach is an assertion that, under certain conditions, a probability can be ascribed to a complete history of a quantum system without invoking any external state-vector reductions in the development of the history. Any such scheme would clearly be particularly attractive in quantum cosmology where a fundamental observer-system split seems to be singularly inappropriate.

Dowker and Kent 1 have raised some serious doubts in the context of their penetrating analysis. [365], see also L Smolin [13].

the possibility that the decoherent histories programme could provide a framework for solving certain technical or structural problems that arise in quantum gravity. An example is the problem of time that features prominently in canonical quantum gravity (see [364]).

• **decoupling:** this is the instant in the history of the universe where matter (baryons

and electrons) and the electromagnetic radiation cease to be coupled. After this point . This took place when the temperature was about 4000 K.

- **deformed special relativity:**

Deformed special relativity provides an arena in which to build an effective theory for quantum gravity. [256].

- **density matrix:** mixed state described by a density matrix  $\rho$ . Expectation values of observables are given by

$$\rho[A] = \text{tr}[A\rho]. \tag{A.27}$$

- **de Sitter universe:** The de Sitter universe plays an important role in cosmology as it can be used as a good approximation for the stage of accelerated expansion - inflation.

- **DeWitt, Bryce:** Made major contributions to classical and quantum field theory, in particular, to the theory of gravitation. Through application of Dirac's theory of constraints in development of a canonical approach to quantum gravity, DeWitt was led to what became to be known as the Wheeler-DeWitt equation (the Hamilton constraint in 'metric variables'), which has since been applied many times to problems in quantum cosmology.

By the end of 1965 he had found the rules for quantising the gravitational and non-Abelian gauge fields to all orders nearly two years before a paper by Fad'eev and Popov deriving the same rule was published.

See BRYCE SELIGMAN DEWITT 1923-2004 - A Biographical Memoir by STEVEN WEINBERG.

- **Dirac conjecture:** All first class constraints generate gauge transformations at a given time.

- **Dirac, Paul:** Developed the hamiltonian formulation of GR independently of Bergmann and his group, with the long term goal to quantize gravity. The main tool for this, the hamiltonian theory of constrained systems, was developed for this purpose.

- **Dirac-Bergman algorithm:** Procedure for generating a finite set of constraints consistent under evolution.

- **Dirac observable:** An observable is a function on the constraint surface that is invariant under gauge transformations generated by *all* the first class constraints. Equivalently, an observable is a function on the phase space that has weakly vanishing Poisson brackets with the first class constraints.

It is often stated that there are no Dirac observables known for General Relativity, except for the ten Poincaré charges at spatial infinity in situations with asymptotically flat boundary conditions.




Figure A.11: Bryce DeWitt.

- **dispersion relations:** The form of the dispersion relations is dictated by Lorentz invariance.
- **Dittrich, Bianca:**
- **domain of dependence:** The future domain of dependence,  $D^+(\Sigma)$ , is the set of points  $p$  in  $\mathcal{M}$  for which every past-inextendable causal curve through  $p$  intersects  $\Sigma$ .
- **Doplicher, Haag, Roberts DHR:** for classification of available representations in order to know which features are tied to a specific representation and which are not.
- **double harmonic oscillator model:** The double harmonic oscillator was first studied by Rovelli [277] as a toy model to help understand the “problem of time”.

The Hamiltonian for the double harmonic oscillator

$$\mathcal{H} = \lambda \left( \frac{1}{2}(p_1^2 + \omega^2 q_1^2) + \frac{1}{2}(p_2^2 + \omega^2 q_2^2) - E \right) \quad (\text{A.28})$$

One of the oscillators can be thought of as a “clock” for the other oscillator.

Other toy models that share some essential features of GR are .

- **doubly special relativity:** Doubly special relativity theories are relativistic theories in which the transformations between inertial observers are characterized by two observer-independent scales - the light speed and the Planck length  $l_p$ . The Planck length seems to have a crucial role to play in quantum theories of gravity as a threshold to quantum



Figure A.12: Paul Dirac.

effects of spacetime. It is argued that the planck length must have the same value in all inertial frames which is in contradiction with special relativity. Double special relativity is a proposal to solve this perceived problem.

A phenomenological description of the signatures of Lorentz invariance violation could be doubly special relativity (DSR) [??], a theory in which not only the speed of light but also the Planck energy is (inertial) frame independent.

- **Dreyer, Olaf:** [19] “...argues that the incompatibility between quantum theory and general relativity can be resolved only if we give up the idea that space is fundamental. He proposes that space itself emerges from a more fundamental description that is quite different.”

- **dust:** Matter that does not interact on one another by mechanical forces but interacts with itself only via the gravitational field.

- **dynamical horizons:** Isolated horizons provide a quasi-local description of black holes in equilibrium. There has also been developed a framework for the quasi-local treatment of black holes that are not in equilibrium (because of infalling matter or interaction with external bodies), namely *dynamical horizons*.

The formal definition of a dynamical horizon is as follows,

- i)  $S$ , the explanation  $\Theta_{(\ell)}$  of the one null normal,  $\ell^a$ , is everywhere zero and that
- ii) that of the other null normal,  $n^a$ , is negative.

The second condition merely says that  $n^a$  is inward pointing null normal. none of the



Figure A.13: Bianca Dittrich.




Figure A.14: The future domain of dependence,  $D^+(\Sigma)$ , of  $\Sigma$ .  $p$  is in  $D^+(\Sigma)$ ,  $q$  isn't because there are past-inextendable causal curve through  $q$  that don't intersect  $\Sigma$ , e.g. the curve  $\gamma$ .

light rays emerging from *any* point on  $S$  are directed towards the 'outside' region.

how black-holes grow

- **dynamical symmetry:** e.g. invariance under active diffeomorphisms in GR.
- **dynamically triangulated quantum gravity:** dynamical triangulations which edge lengths are fixed and the path integral is represented as a sum over triangulation.

A weak coupling phase with quantum fluctuations around a "semiclassical" dynamically generated background geometry despite the fact that the formulation is explicitly background-independent, [264].

- **edge:** Let  $S$  be a closed achronal set. The edge of  $S$  is defined as a set of points  $x \in S$  such that every neighbourhood of  $x$  contains  $y \in I^+(S)$  and  $z \in I^-(S)$  with a timelike

curve from  $z$  to  $y$  which does not meet  $S$ .

Link between vertices of a graph.

• **Ehrenfest's theorem:**

$$\frac{d\langle\hat{x}\rangle}{dt} = \frac{\langle\hat{p}_x\rangle}{m}, \quad (\text{A.29})$$

$$\frac{d\langle\hat{p}_x\rangle}{dt} = -\left\langle\frac{\partial\hat{V}}{\partial x}\right\rangle. \quad (\text{A.30})$$

$$\frac{d\hat{x}}{dt} = \frac{i}{\hbar}[\hat{\mathcal{H}}, \hat{x}], \quad \frac{d\hat{p}_x}{dt} = \frac{i}{\hbar}[\hat{\mathcal{H}}, \hat{p}_x] \quad (\text{A.31})$$

$$\hat{\mathcal{H}} = \frac{1}{2m}(\hat{p}_x^2 + \hat{p}_y^2 + \hat{p}_z^2) + \hat{V}(x, y, z) \quad (\text{A.32})$$

$$\frac{d\hat{x}}{dt} = \frac{\hat{p}_x}{m}, \quad \frac{d\hat{p}_x}{dt} = -\frac{\partial\hat{V}}{\partial x} \quad (\text{A.33})$$

$$\frac{d}{dt} \int \psi^* \hat{x} \psi dx = \frac{1}{m} \int \psi^* \hat{p}_x \psi dx, \quad \frac{d}{dt} \int \psi^* \hat{p}_x \psi dx = - \int \psi^* \frac{\partial\hat{V}}{\partial x} \psi dx \quad (\text{A.34})$$

• **Einstein, Albert-(1878-1955):** Began his scientific work at the beginning of the 20th century. During the years 1905-1906 he published three articles: (1) The special theory of relativity; (2) an explanation of the photo-electric effect; (3) Brownian motion. The three corner stones of modern physics. Considered to be very clever.

• **Einstein's hole argument:** As explained by Rovelli “Assume the gravitational field equations are covariant. Consider a solution of these equations in which the gravitational field is  $e$  [basically the metric] and there is a region  $H$  of the universe without matter (the “hole”). Assume that inside  $H$  there is a point  $A$  where  $e$  is flat and the point  $B$  where it is not flat. Consider a smooth map  $\phi: \mathcal{M} \rightarrow \mathcal{M}$  which reduces to the identity outside  $H$ , and such that  $\phi(P) = Q$ , and let  $\tilde{e} = \phi_* e$  be the pull back of  $e$  under  $\phi$  [the metric induced by the diff transformation  $\phi$ ]. Notice that  $e$  is flat around the event  $A$  while  $\tilde{e}$  is not. The two fields  $e$  and  $\tilde{e}$  have the same past, are both solutions of the field equations, but have different properties at the event  $A$ , therefore they are not deterministic. But we know that (classical) gravitational physics is deterministic, therefore, either the field equations can or cannot be generally covariant, or there is no meaning to the physical event  $A$ .”

• **eikonal approximation:** an approximation for the scattering amplitude valid in the high energy, low momentum transfer regime,




Figure A.15: Albert Einstein (1879-1955).

$$\mathbf{p}' = \mathbf{p} - \frac{\mathbf{k}}{2}, \quad \mathbf{p}'' = \mathbf{p} + \frac{\mathbf{k}}{2}, \quad \mathbf{p}^2 \rightarrow \infty, \quad \mathbf{p}^2 > \mathbf{k}^2 \quad (\text{A.35})$$

The dominant contributions come from classical trajectories.

- **eikonal equation:**
- **electroweak theory:** combining theory of QED and weak-interactions.
- **energy conditions:**

**The weak energy condition:**

$$T^{ab}u_a u_b \geq 0, \quad (\text{A.36})$$

for all timelike vectors  $u^a$ . This condition requires that the local energy density be non-negative in every observer's frame. Again this seems to be a very reasonable condition in classical theory.

**The strong energy condition:**

$$((T^{ab} - \frac{1}{2}g^{ab}T)u_a u_b \geq 0, \tag{A.37}$$

for all timelike vectors  $u^a$ . The frame in which  $T^{ab}$  is diagonal, this condition implies that the local energy density  $\rho$  plus the sum of local pressures  $p^i$  is non-negative:  $\rho + \sum_i p^i \geq 0$ , and that  $\rho + p^i$  for each  $i$ . This condition certainly holds for ordinary forms of matter, although it can be violated of classical physics.

**The null energy condition:**

$$T^{ab}k_a k_b \geq 0, \tag{A.38}$$

This condition is implicit in the weak energy condition. That is, if we assume the weak energy condition, then the null energy condition follows by continuity as  $u^a$  approaches a null vector.

**The dominant energy condition:**

The dominant energy condition implies that matter cannot travel faster than the speed of light.

$$R_{ab}W^a W^b \geq 0 \tag{A.39}$$

for any null vector  $\mathbf{W}$ . The importance of the weak energy condition is that it implies that in presence of matter a congruence of null geodesics that are initially diverging will begin to .

$$\frac{d}{d\lambda}\hat{\theta} = -R_{ab}K^a K^b - 2\hat{\sigma}^2 - \frac{1}{2}\hat{\theta}^2 \leq 0. \tag{A.40}$$

$\hat{\theta}$  monotonically decreases along the null geodesics if the first term on the right hand side if the weak energy condition holds.

- **energy conservation:** Notion of energy and the law of energy conservation have played a key role in analyzing behaviour of physical theories. However, it is not an independent physical requirement, but rather a consequence of time translation symmetry of the background metric and the dynamical equations.

Energy conservation is just a consequence of invariance under time translations, which, it turn is a feature of the homogeneity of Minkowski spacetime. However, when we come to general relativity the homogeneity of the Minkowski solution is not feature invariant under active diffeomorphisms. Hence there is no fundamental energy conservation in nature - there is nothing sacred about energy conservation!

When we restrict possible spacetimes to be those that are asymptotically flat, diffeomorphisms at infinity are just those of time and space translation we recover our notions of energy and momentum conservation.

In the absence of fixed a spacetime we must learn how to do physics in the absence of energy or momentum.

Define  $E = v^a K_a$ , where  $K_a$  is a killing vector,

$$\begin{aligned}\nabla_v E &= v^a \nabla_a (v^c K_c) \\ &= v^a v^c \nabla_a K_c \quad \text{since } v \text{ is the tangent of a geodesic,} \\ &= v^a v^c \nabla_{(a} K_{c)} = 0\end{aligned}\tag{A.41}$$

- **Euclidean geometry:** Flat geometry based upon the geometries axioms of Euclid.
- **energy momentum tensor  $T_{ab}$ :**
- **entropy:** A quantitative measure of disorder of a system. The greater the disorder, the higher the entropy. It is defined as the amount of information about the microscopic motion of the constituents making up the system which are not determined by a description of the macroscopic state of that system.
- **epistemology:** Philosophic subject concerned with the significance of what we can know.
- **EPR:**

Rovelli's book [20]

*The EPR apparent paradox might be among these. The two observers far from each other are physical systems. The standard account neglects the fact that each of the two is in a quantum superposition with respect to the other, until the moment they physically communicate. But this communication is a physical interaction and must be strictly consistent with causality.*

- **entanglement entropy:** Black hole entanglement entropy is a measure of the information loss due to separation between

inside and outside the event horizon.

- **equations of motion (EOM):** They are a prescription for calculating the acceleration from the state of the system (from initial position and velocity). The initial state yields the initial acceleration, and the integration leads from the initial acceleration to later sates.

- **event:** A point in four-dimensional: a location in both space and time.

- **Event horizon:** A surface that divides spacetime into two regions: that which can be observed and that which can not. The Schwartzschild radius of a nonrotating black hole is an event horizon.????????

- **evolving Constants of motion:** Introduced by roveli in [279] (an idea that goes back to DeWitt, Bergmann nad Einstein). The Hamiltonian constraints generate the dynamics, dynamics is pure gauge, and the observables of the theory are constant along the gauge orbits - these are the evolving constants of the motion. They are formed as follows: in a totally constrained theory, the values of fields are not physically observable. On the other hand, if one chooses a one-parameter family of observables such that their value coincides with the value of a dynamical variable when the parameter takes the value of another dynamical variable, which one uses to characterize the evolution.

- **exceptional edges:** Exceptional edges are characterized by not being effected by action of the Scalar constraint. gr-qc/0409061

Application of the Hamiltonian constraint on a vertex  $v$  will produce a new edge  $a_{12}$  fig.(A.16) attached to (two) the edges  $e_1$  and  $e_2$ . The edges incident to the vertices  $v_1$  and  $v_2$  only have two linearly independent tangent directions, namely those of  $e_1$  and  $a_{12}$ . An exceptional edge, here  $a_{12}$ , verticies are annihilated by a second application of the Hamiltonian constraint. A failure to ‘propagate’.




Figure A.16: exceptional. The Hamiltonian constraint’s action on the trivalent node  $v$  with linearly independent tangent vectors produces two new vertices  $v_1$  and  $v_2$  - each of which has only two linearly independent tangent vectors.

- **Extended Master Constraint Operator  $\hat{M}_E$ :**

$$\hat{M}_E := \int_{\sigma} d^3x \frac{C^2 + q^{ab}C_a C_b}{\sqrt{\det(q)}} \tag{A.42}$$

Both constraints are spatially diffeomorphism invariant. However,  $\hat{M}_E$  allows us to implement both the Hamiltonian and the spatial diffeomorphism constraint on  $\mathcal{D}_{kin}$  provided we implement the corresponding operators in a non graph changing fashion. In [77] it was shown how to do that using the notion of a minimal loop which is a loop (average) within the graph on which the constraint acts. It follows that instead of using dual operators we

can directly work with operators on  $\mathcal{H}_{kin}$  and their adjoints so that the construction of the quadratic form can be sidestepped.

There is the Extended Master Constraint Operator  $\hat{M}_{EE}$  which also includes the Gauss constraint in addition:

$$\hat{M}_{EE} := \int_{\sigma} d^3x \frac{C^2 + q^{ab} C_a C_b + C_j C_j}{\sqrt{\det(q)}}. \quad (\text{A.43})$$

See page ?? of the Feynman lectures.

• **extensive thermodynamical quantities:**

• **extremal black holes** Black holes having the maximum possible charge for a given mass and angular momentum. Thought to be astronomically extremely rare. Ruled out by the third law of black hole mechanics.

• **extrinsic curvature:** We can describe this curvature by looking at how the normal vectors to the surface change along the surface. One uses the derivative operator of the larger space to measure the changes in the normal vector field, and then project back into the surface.

extrinsic curvature of spacetime  $\mathcal{M}$  to measure the changes in the timelike normal vector field of the slice  $\Sigma$ . Roughly speaking, it is to do with the time derivative of the 3-metric.

Contains the information on how a hypersurface is embedded in the enveloping spacetime.

• **faithful:**

**faithful representation:**

**faithful state:**

• **false vacuum:** A metastable state in which a quantum field is zero, but its corresponding vacuum energy density is not zero.

• **falsifiable:** The property of a scientific hypothesis that it is not possible to perform an experiment that would disprove, or falsify, the hypothesis.

• **Fell's theorem:** (From [11] Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics, Robert M. Wald) For an algebra of observables  $\mathcal{A}$  of a field theory, although two representations of  $\mathcal{A}$  may be unitarily inequivalent, the determination of a finite number of expectation values of observables in  $\mathcal{A}$  - each made with finite accuracy - cannot distinguish between different representations.

Quote:

whether the resulting physical Hilbert space admits a sufficient number of physical semi-classical states - in some sense this is granted by Frell's theorem [114] once we have a representation of the physical Hilbert space of the C\* algebra of our preferred algebra of bounded Dirac observables. [208]

- **Fermions:** The metric formulation cannot incorporate fermionic equations of motion. The reason for this is there is no finite-dimensional spinorial representation of the  $4 \times 4$  general-linear real matrix group, however, there is a finite-dimensional spinorial representation of the Lorentz group - which is what Dirac discovered. We use a tetrad frame to establish an instantaneous inertial frame, in which the laws of special relativity apply. It can be introduced in a generally covariant way. When we fix the tetrad to Minkowski spacetime we recover Dirac's equation.

- **Fermi-Walker derivative:**

- **Fermi-Walker propagated:** propagation along a non-geodesic curve.

- **Feynmann diagrams:**




Figure A.17: The charge renormalization.

Feynman graphs yield integrals that diverge when momentum becomes indefinitely large (or, equivalently, when distances become indefinitely small).

one-particle irreducible (1PI) diagrams

- **Feynmann-Kac formula:** Instead of considering matrix elements of the unitary operator  $U(t) = \exp(itH/\hbar)$ , consider the analytic continuation  $S^{A_1 \dots A_n}(t_1, \dots, t_n)$ , of the Wightman functions  $W^{A_1 \dots A_n}(t_1, \dots, t_n)$  is

$$S^{A_1 \dots A_n}(t_1, \dots, t_n) := W^{A_1 \dots A_n}(t_1, \dots, t_n).$$

These are the so-called Schwinger functions. A path integral for the Schwinger functions have better chance to be rigorously defined because the measure has a damping factor rather than an oscillating one and so one can give a rigorous definition by backwards analytic continuation (when possible). The path integral for the Schwinger functions (when it can be proved) is called the Feynmann-Kac formula.

- **first class constraints:** restricting the fields to the constraint surface.

infinitesimal “gauge transformation”

$$(1 + a \frac{\partial}{\partial x})f(x) = f(x + a) + \mathcal{O}(a^2) \quad (\text{A.44})$$

Functions invariant under this gauge transformation satisfy the constraint equation,

$$\frac{d}{dx}f(x) = 0. \quad (\text{A.45})$$

The first class property is that

$$\{C_m, C_n\} = C_{mn}{}^l C_l \quad (\text{A.46})$$

where the  $C_{mn}{}^l$  are called the structure constants if they are independent of on the phase space coordinates and structure functions if they are functions over phase space; that is the first class constraints satisfy, in general, a “non-Lie” algebra

$$C_{mn}{}^l = C_{mn}{}^l(q^i, p_i) \quad (\text{A.47})$$

as it is possible that from the canonical variables.

The first class property of the constraints guarantees, that the flow of the constraints is integrable to an  $n$  dimensional surface - the gauge orbit.

they generate gauge transformations.

in the Hamiltonian approach based on Dirac’s conjecture [72], where a suitable combination of the first class constraints is shown to to be a generator of local symmetries of the Lagrangian.

as apposed to a structure *constant* we can have a structure *function* depending on the phase space coordinates.

**Dirac’s conjecture** in the Hamiltonian approach the symmetries are generated by the first class constraints.

In the Lagrangean approach local symmetries are reflected in the existence of so called gauge identities.

- **first class constraints:** When all the constraints are first class we then have a closed constraint algebra.

- **flatness problem:** The observed fact that the geometry of of the universe is nearly very nearly flat, a very special condition, without an explanation of why it should be.
- **Fleischhack, Christian:**



Figure A.18: Christian Fleischhack.

- **Fock representation:** The most commonly used representations of canonical commutation relations/canonical anti-commutation relations are the so called Fock representations, defined in bosonic/fermionic Fock spaces. These spaces have distinguished vector  $\Omega$  called the vacuum killed by the annihilation operators and cyclic with respect to creation operators (i.e., repeated applications and complex linear combinations of creation operators on the  $\Omega$  generates a dense subset of the Hilbert space). They were introduced to describe systems of many particle systems with Bose/Fermi statistics.

Fock space is considered the prototypical QFT state space by some

fundamentally the Fock representation is not a valid representation for interacting quantum field theories.

States in LQG are fundamentally different from Fock states of Minkowskian quantum field theories. The main reason is the underlying diffeomorphism invariance: In the absence of a background geometry, it is not possible to introduce the familiar Gaussian measures and associated Fock spaces.

- **foliation of space time:** We have a spacial three-space stacked one above the other, one for every “moment in time”.
- **form factor:** Given a loop  $\gamma$ , its form factor  $F^a(\gamma, \vec{x})$  is defined by

$$F^a(\gamma, \vec{x}) := \oint_{\gamma} ds \dot{\gamma}^a \delta^3(\vec{x}, \vec{\gamma}(s)) \quad (\text{A.48})$$

- **focussing theorem:** It is easy to show that the Raychaudhuri equation implies the following inequality for the expansion  $\theta(\lambda)$

$$\theta^{-1}(\lambda) \geq \theta^{-1}(0) + \frac{\lambda}{4} \tag{A.49}$$

From this inequality we see that if the congruence is initially converging ( $\theta(0) < 0$ ), then  $\theta(\lambda) \rightarrow -\infty$  within an affine parameter  $\lambda \leq 2/|\theta(0)|$ . Demonstrates attractive instability of gravity. This result is employed in the area increase law of black holes and in the singularity theorems.

- **frame of reference:** The coordinate system to which a particular observer refers his or her measurements.

- **free-fall:** a particle is said to be in free-fall when its motion is affected by no forces except gravity. Unrestrained motion under the influence of the gravitational field.

or

A body in free fall does not feel any acceleration, and hence behaves locally as an inertial reference system of SR.

It is meaningless to say whether or not the gravitational field is flat around some point  $p$  of  $\mathcal{M}$ . It is meaningful, however, to ask whether or not the gravitational field is flat around the place where two particles coincide.

- **free data:** isolated horizons. reconstruct from...

- **Friedmann models:** A class of cosmological models that are isotropic and homogeneous, contain specified matter-energy density, conserve matter, and admit no cosmological constant. Also called standard models.

[307]

All textbooks on classical GR incorrectly describe the Friedmann equations as physical evolution equations rather than what they really are, namely gauge transformation equations. The true evolution equations acquire possibly observable modifications to the gauge transformation equations whose magnitude depends on the physical clock that one uses to deparametrise the gauge transformation equations.

- **functional analysis:** is the branch of mathematics which is concerned with the study of spaces of functions.

- **functional integration:**

In the Hilbert space language, this is the origin of field theoretic infinities, e.g., the reason why we can not naively multiply field operators.

- **future-distinguishing spacetime:** A spacetime is said to be future-distinguishing if  $a \neq b$  implies  $I^+(a) \neq I^+(b)$ . (past-distinguishing is similarly defined: if  $a \neq b$  implies  $I^-(a) \neq I^-(b)$ ). (See also chronology condition, strongly causality, stable causality).




Figure A.19:

- **Gambini, Rodolfo:**



Figure A.20: Rodolfo Gambini.

- **gamma ray burst:** A  $\gamma$ -ray burst is a light signal of extremely high energetic photons (up to 1 TeV) that travels over cosmological distances ( $\approx 10^9$  years). Could have applications in quantum gravity phenomenology.

They are enormous explosions that for a few seconds can produce as much light as that emitted by a whole galaxy. Signals from these explosions reach Earth on average once about once a day.

- **gauge fixing:** One method to eliminate the superfluous degrees of freedom of a gauge theory is through the introduction of a new set of constraints.

leads to second class constraints. null components? Poisson bracket is replaced by the Dirac bracket

$$\{f, g\}^* := \{f, g\} - \{f, C_i\} B_{ij}^{-1} \{C_j, g\} \quad (\text{A.50})$$

where

$$B_{ij} := \{C_i, C_j\} \quad (\text{A.51})$$

The geometric interpretation of this is: the symplectic structure induced on the hypersurface, defined by the condition  $C_j = 0$ ?, from the symplectic structure of the phase space  $\mathcal{M}$  is given by the Dirac bracket  $\{\cdot, \cdot\}^*$ .

Interpretation of gauge invariant variables is based on the coincidence of the variables in an appropriate gauge choice. For example,  $\text{blah}$  defined by  $\text{blah}$  describes  $\text{blah}$ , because the variable  $\text{blah}$  coincides with  $\text{blah}$ , defined in  $\text{blah}$  in the gauge choice  $\text{blah}$ .

- **gauge invariance:**
- **gauge invariant coherent states:** Projections of the complexifier coherent states onto the gauge-invariant subspace, [253], [254].
- **gauge potential:** Gauge potentials can be identified as the connection on a principal fibre bundle in a certain gauge.
- **gauge symmetry:**

A gauge symmetry is a redundancy in the description of the system and not a symmetry of a system. For example in a situation with rotational symmetry, such as a planet moving about the sun, although there is a symmetry that changes the angle in the plane of motion there is still physically a difference between two different values of the angular variable. In a gauge theory this is not the case. If two field configurations are related by a gauge transformation then they are physically identical.

Dirac's definition of a gauge symmetry is when two solutions with the same initial data set separate at some point.

- **gauge transformations:** Gauge transformations are maps from solutions of the equations of motion into other solutions which are physically equivalent.

There is a choice of a gauge potential in a certain chart but there is also a choice of local charts. This other flexibility means that a gauge does not just refer to a particular gauge potential.

In the physics literature gauge transformations often refer to passing from one chart to another in a overlapping region - a passive gauge transformation. In the physics literature gauge transformations always refer to passing from one gauge potential to a physically equivalent one in the same chart - an active gauge transformation. It is active gauge transformations that have physical meaning.

electromagnetism the gauge transformation of the gauge potential is

$$\delta A_\mu = \{A_\mu, G\} = \partial_\mu \Lambda, \quad (\text{A.52})$$

for an arbitrary function  $\Lambda$ .

In general relativity the gauge transformations (diffeomorphisms) for the metric field

$$\delta g_{\mu\nu} = \epsilon^\rho \partial_\rho + g_{\mu\rho} \partial_\nu \epsilon^\rho + g_{\rho\nu} \epsilon^\rho, \quad (\text{A.53})$$

for some arbitrary function  $\epsilon^\rho$ .

• **Gauss-Codaza equations:** Gauss-Codacci equation relating the space-time curvature to the intrinsic curvature of a sub-manifold. The relation can be written:

$$R_{qrt}^p = \Gamma_{qr|t}^p - \Gamma_{qt|r}^p + \Gamma_{qr}^s \Gamma_{st}^p - \Gamma_{qt}^s \Gamma_{sr}^p. \quad (\text{A.54})$$

See maths glossary.

This is not the same as saying that “General covariance is the idea that the laws of nature must be the same in all reference frames, and hence all coordinate systems”.

**Gelfand triple:** In LQG the nice quantum states are typically taken to be finite linear combinations of spin network states whose space is denoted by  $Cyl^*$  (the space of cylindrical functions of connections). Solutions to the quantum Einstein equations belong to its dual,  $Cly^*$ .

• **generating functional:**

$$W[J] = \int \mathcal{D}\phi \exp\left(\frac{i}{\hbar} \int_{-\infty}^{\infty} [\mathcal{L}(\phi, \partial_\mu \phi) + J\phi] d\tau\right) \quad (\text{A.55})$$

$$N^{-1} \exp\left(i \int \int \mathcal{L}[\phi]\right) \prod_x [d\phi(x)] \quad (\text{A.56})$$

in QFT the generating functional (A.55) is introduced as formal extrapolation of measures in infinite-dimensional spaces

$$\prod_x [d\phi(x)]. \quad (\text{A.57})$$

This fails to be a true measure because the Lebesgue measure in infinite-dimensional spaces fails to be defined in general.

Given a connection  $A$  all its gauge invariant information is contained in all graphs with a finite number of degrees of freedom. We have a genuine field theory with an infinite number of degrees of freedom.

called the vertical polarization.)

- **general boundary formulation:** The so-called *general boundary formalism* extends the usual formulation of dynamics in terms of space-like hypersurfaces to one with closed boundary surfaces.

Homepage of Robert Oeckl <http://www.matmor.unam.mx/~robert/>

*“Quantum gravity is the search for a theory that may unify general relativity with quantum mechanics and ultimately with the standard model of elementary particle physics. This problem has motivated me to think about the foundations of quantum theory. In particular, I am trying to extend the standard formulation of quantum mechanics in such a way as to make it more compatible with general relativistic concepts. It turns out that this is best done directly in the context of quantum field theory rather than non-relativistic quantum mechanics. I call this extension tentatively the general boundary formulation.”*

- **general covariance:** The principle that the laws of nature must be the same in all reference frames. From [17] (p.469): there should not be special coordinates that have a physical role to play, and that the equations of the theory should be such that their most natural expression does not depend on any particular choice of coordinates.

- **general relativity:** Often described as a theory that conceives gravity not as a force between masses but as a change in the geometry of spacetime due to the presence of the matter and energy. However, the full content of GR is that it *discards the very notion of spacetime*. Instead, “spacetime geometry” is conceived as certain aspects of the relationships that exist between physical objects that live in the world (this what general relativists are referring to when they say that GR is a *background independent* theory). This is the context for Einstein’s remark “*Beyond [my] wildest expectations*” (see Einstein’s hole argument, active diffeomorphisms)

- **generalized fields:** in the passage from the classical field to the quantum field. with classical fields there is no classical particle picture space of distributional fields - smooth classical fields have measure zero in the quantum configuration space.

In quantizing a finite dimensional system one defines states as functions on a configuration space and then defines an inner product of two such functions  $\psi$  and  $\phi$  through

$$(\psi, \phi) = \int_{\mathcal{Q}} d\mu \psi^* \phi \tag{A.58}$$

In the quantum theory systems with a finite number of degrees of freedom - the classical and quantum configuration spaces agree. In field theory the quantum configuration

space is a substantial enlargement of the classical; in scalar field theories, for example, although the classical configurations are smooth functions on a  $t = \text{const}$  slice, the quantum configuration space consists of all tempered distributions.

The set of smooth configurations is of zero measure with respect to the Gaussian measure that determines the inner product! That is, for two smooth wavefunctions  $\Psi$  and  $\Phi$

$$\int_{\mathcal{Q}} d\mu \Psi^* \Phi = 0 \tag{A.59}$$

The regularization problems of quantum field theory can be traced back to this fact.

It is the distributional field fluctuations at the horizon that are the microscopic source for black hole entropy.

- **generalized Wick transformation:** which is arising by relating simpler Riemannian quantum theory to the more complicated physical Lorentzian theory. Allows one to obtain physical states in Lorentzian gravity from Wick transforming solutions of Riemannian quantum theory.

- **generic condition:**

(i) The strong energy condition holds,

(ii) Every timelike or null geodesic contains a point where

$$\ell_{[a} R_{b]dc[e} \ell_{f]} \ell^c \ell^d \neq 0, \tag{A.60}$$

( $\ell_a$  being the tangent vector of the geodesic). The condition serves to exclude certain pathological spacetimes ... I think?

see energy conditions - strong and weak.

- **geodesic:** A geodesic is the closest thing there is to a straight line curved space time. The shortest distance between two points. A more general definition of a geodesic is its velocity vector is parallel transported along the curve it traces out in spacetime. In other words, the parallelly propagated vector at any point of the curve is parallel, that is, proportional to the vector at this point:

$$\frac{d^2 x^a}{du^2} + \Gamma_{bc}^a \frac{dx^b}{du} \frac{dx^c}{du} = \lambda \frac{dx^a}{du}. \tag{A.61}$$

- **geodesically convex:** A set  $U$  of spacetime  $\mathcal{M}$  is geodesically convex if any two points of  $U$  may be joined by a unique geodesic lying entirely in  $U$ .

- **geometric quantization:** Geometric quantization provides a geometric, general framework for the quantization of a symplectic manifold  $M$  which is not necessarily a cotangent bundle, for example when  $M$  is compact.

arXiv:quant-ph/0006037 v1 7 Jun 2000 HARMONIC ANALYSIS WITH RESPECT TO HEAT KERNEL

Roughly a polarization is a choice of  $d$  coordinates on the  $2d$ -phase space  $\mathcal{M}$ , with the idea that the functions in our quantum Hilbert space will be independent of these  $d$  variables.

For example if  $\mathcal{M} = \mathbb{R}^{2d}$ , then we may take the usual position and momentum variables  $x_1, \dots, x_d, p_1, \dots, p_d$  and then consider functions that depend only on  $x_1, \dots, x_d$  and are independent of  $p_1, \dots, p_d$ . (This is called the vertical polarization.)

Alternatively, one may consider complex variables  $z_1, \dots, z_d, \bar{z}_1, \dots, \bar{z}_d$  and then consider the functions that are independent of  $\bar{z}_k$ , that is, holomorphic. (This is a complex polarization.) In that case our Hilbert space is the Segal-Bargmann space. To be more precise, in geometric quantization the elements of the quantum Hilbert space are not functions, but rather sections of a certain complex line bundle with connection. The sections are required to be covariantly constant in the directions corresponding to the polarization.

Apparent dependence of the act of quantization on the choice of coordinates.

- **geons:**

Lines of force can disappear at one point and reappear at the other - one point acts as sink and the other a source. The field equations of gravity would determine the motion of the sources and sinks and therefore how the charged particles represented by them. Purely gravitational effects.




Figure A.21: geons.

- **Gibbs state:**

$$\exp(\mathcal{H}/kT) \tag{A.62}$$

- **Gleason’s theorem:**

- **globally hyperbolic:** Globally hyperbolicity is a property of some spacetimes with the topology  $\mathbf{R} \times \Sigma$ , it ensures that there exists a universal time function whose gradient is everywhere timelike.

The physical significance of global hyperbolicity comes from the fact that there is a family of Cauchy surfaces  $\Sigma(t)$  for a region of spacetime  $\mathcal{M}$ . One can predict what will happen in from data on the Cauchy surface.

In a globally hyperbolic region of spacetime, there is a geodesic of maximum length joining any pair of points that can be joined by a timelike or null curve. This fact is used in the singularity theorems: one establishes there is a region of spacetime which is globally hyperbolic, then one demonstrates that there is a geodesic which has a pair of conjugate points and hence is not a of doesn’t maximum length (involves focussing theorem). The way out of this contradiction is that this geodesic comes to a full stop indicating the presence of a singularity.

Equivalent definition of global hyperbolicity: A spacetime  $\mathcal{M}$  is said to be globally hyperbolic if the sets  $J^+(x) \cap J^-(y)$  are compact for all  $x, y$  in  $\mathcal{M}$

The global hyperbolicity of  $\mathcal{M}$  is closely related to the future or past development of initial data from a given spacelike hypersurface.

- **“global time problem”**

“multiple choice problem” - that one could use different time functions and one must prove the quantum theories are equivalent.

[?] Arlen Anderson, *Evolving Constants of Motion*, [gr-qc/95070708].

- **Gowdy spacetime:** an example of a that can be quantized as a midisuoerspace model.

- **gravitational backgrounds:**

$$\mathcal{L} = \sqrt{-g}\{g^{\mu\nu}\nabla_\mu\phi^*\nabla_\nu\phi - m^2\phi^*\phi\} \tag{A.63}$$

$$g^{\mu\nu}\nabla_\mu\nabla_\nu\phi + m^2\phi = 0 \tag{A.64}$$

and the conserved current density is:

$$J_\mu = i(\phi^*\nabla_\mu\phi - (\nabla_\mu\phi^*)\phi) \equiv i\phi^*\overleftrightarrow{\nabla}_\mu\phi \tag{A.65}$$

the inner product:

$$\langle \psi | \phi \rangle = i \int_{\Sigma} \sqrt{-g} \psi^* \overleftrightarrow{\nabla}_{\mu} \phi d\Sigma^{\mu} \quad (\text{A.66})$$

is independent of  $\Sigma$ .

- **“gravitational bound”**: the gravitational bound problem in GR is an intrinsically non-linear problem even when the conditions are such that the field is weak and the motions are non-relativistic, at least in the time-dependent case. [?]
- **gravitational wave**: A gravitational wave is a linear perturbation of the gravitational field characterised by an oscillating curvature tensor which causes the immediate neighbourhood of space-time region through which it passes to oscillate.
- **gravitons**: Gravitons don't play a fundamental role in the loop quantum gravity. Instead, they only serve as a kind of approximate way of dealing with small perturbations of a wave state in which a superposition of spin networks mimic a given classical spacetime metric.

*“There is a general belief, reinforced by statements in standard textbooks, that: (i) one can obtain the full non-linear Einstein's theory of gravity by coupling a massless, spin-2 field  $h_{ab}$  self-consistently to the total energy momentum tensor, including its own; (ii) this procedure is unique and leads to Einstein-Hilbert action and (iii) it only uses standard concepts in Lorentz invariant field theory and does not involve any geometrical assumptions. After providing several reasons why such beliefs are suspect and critically re-examining several previous attempts we provide a detailed analysis aimed at clarifying the situation. First, we prove that it is impossible to obtain the Einstein-Hilbert (EH) action, starting from the standard action for gravitons in linear theory and iterating repeatedly. Second, we use the Taylor series expansion of the action for Einstein's theory, to identify the tensor  $\mathcal{S}^{ab}$ , to which the graviton field  $h_{ab}$  couples to the lowest order. We show that the second rank tensor  $\mathcal{S}^{ab}$  is not the conventional energy momentum tensor  $T^{ab}$  of the graviton and provide an explanation for this feature. Third, we construct the full nonlinear Einstein's theory with the source being spin-0 field, spin-1 field or relativistic particles by explicitly coupling the spin-2 field to this second rank tensor  $\mathcal{S}^{ab}$  order by order and summing up the infinite series. Finally, we construct the theory obtained by self consistently coupling  $h_{ab}$  to the conventional energy momentum tensor  $T^{ab}$  order by order and show that this does not lead to Einstein's theory. (condensed).” There is more to gravity than gravitons and this will be elaborated in a separate publication, [in preparation].*

Fock states are not available in the absence of background spacetime - so what are particles? ????

- **Groenwald-van Hove theorem**: A collection  $f_i$  of functions on the symplectic space  $(\mathcal{M}, \omega)$  which Poisson commute with each other (are in involution) is said to be complete if the vanishing of  $\{f_i, g\}$  for all  $i$  implies that  $g$  is a function of the form

$g(x) = h(f_1(x), \dots, f_n(x))$ . A collection of operators  $\{A_j\}$  is said to be complete if any operator  $B$  which commutes with each  $A_j$  is a multiple of the identity. Consider a linear map  $\rho$  from the space of observables  $C^\infty(\mathcal{M})$  on the symplectic space  $(\mathcal{M}, \omega)$  such that

(1)  $\rho(1) = \mathbb{I}$ ,

(2)  $\rho(\{f, g\}) = i[\rho(f), \rho(g)]/\hbar$ ,

(3) for some complete set of functions  $f_1, \dots, f_n$  in involution, the operators  $\rho(f_1), \dots, \rho(f_n)$  form a complete commuting set.

The Groenwald-van Hove theorem states that such a map does not exist in general.

- **group averaging:** Projecting it onto the physical Hilbert space by gauge-orbit smearing. We are using the finite - rather than the infinitesimal - form of the constraints.

you have to always first of all you should define a subspace in the Hilbert space which should have stronger topology than the topology of the Hilbert space.

- **Gribov problem:** An obstruction occurs when the gauge fixing term can not be defined globally or that the gauge fixing function intersects a gauge orbit more than once.

- **Haag-Kastler axioms:** [32] [393] [394] Axioms of algebraic quantum field theory.

- **Haag-Ruelle theory:** In standard scattering theory one makes the physical assumption that in the far future  $t_f \rightarrow \infty$  and far past  $t_i \rightarrow -\infty$  any outgoing and ingoing particles respectively do not interact. This is not really true. However, using the methods of local quantum physics, assuming that the theory has a mass gap (the four momentum squared operator should have a pure point spectrum which is separated from the continuum) one can prove that the vacuum correlators of the asymptotic fields reduce to those of free field, where the vacuum really means the interacting vacuum.

- **Haag's theorem:** In quantum field theory, as opposed to quantum mechanics, there exist unitary inequivalent representations (that is, representations which cannot be connected by a unitary transformation) of the canonical commutation relations (CCR). Haag's theorem shows that the representation of the interacting theory differs from that of the free theory. This poses problems for perturbative techniques used in standard textbooks on quantum field theory. The free part is used to define an orthonormal basis of states to which the interaction applies. Haag's theorem says that this is not possible. Haag's theorem forbids us from applying perturbation theory we learned in quantum mechanics to quantum field theory. We are left with the question of why perturbation theory works as well as it does.

- **habitat:** Let  $\mathcal{D}$  be the finite linear span of SNWF's which is dense in the kinematic Hilbert space and let  $\mathcal{D}^*$  denote its algebraic dual. A habitat is a subspace of  $\mathcal{D}^*$  containing  $\mathcal{D}_{Diff}^*$  with the minimal requirement that is preserved by the dual action of the Hamiltonian constraints.

raises questions about the Hamiltonian constraints introduced by Thiemann see [].

habitat, ??.

- **Hamilton constraint:** Hamilton constraint generates “time” translations and hence encodes the dynamics of the theory.

$$\hat{\mathcal{H}}_\epsilon(N)|f\rangle = 0 \tag{A.67}$$

$$\langle \Psi | . \rangle \leftrightarrow \Psi( . )$$

$$\Psi( . ) \in (\mathcal{H}_{Kin}^*)_{Diff} \tag{A.68}$$

$$\langle \Psi | \hat{\mathcal{H}}_\epsilon(N)f \rangle \leftrightarrow \Psi(\hat{\mathcal{H}}_\epsilon(N)f) = 0 \tag{A.69}$$

it is independent of the value of  $\epsilon$ ! The underlying reason is understood to be the diffeomorphism covariance of the graph-dependent triangularization prescription. Proved in section ??? The same reasoning applies to the matter coupled case. The limit  $\epsilon \rightarrow 0$  is therefore already performed; hence the theory, including coupling to the standard model, is manifestly finite and does not require renormalization!

- **Hamilton function:** The Hamilton function of a finite dimensional dynamical system is the value of the action of a solution of the equations of motion, viewed as a function of the initial and final coordinates.

The

gr-qc/0408079

- **Hamilton vector field:** Hamiltonian vector field - Given a manifold  $X$  with a symplectic structure  $\omega$ , any smooth function  $f : X \rightarrow R$  can be thought of as a “Hamiltonian”, meaning physically that we think of it as the energy function and let it give rise to a flow on  $X$  describing the time evolution of states. Mathematically speaking, this flow is generated by a vector field  $v(f)$  called the “Hamiltonian vector field” associated to  $f$

$$\omega(., v(f)) = df \tag{A.70}$$

In other words, for any vector field  $u$  on  $X$  we have

$$\omega(u, v(f)) = df(u) = u f \tag{A.71}$$

The vector field  $v(f)$  is guaranteed to exist by the fact that  $\omega$  is nondegenerate.

- **Hamilton-Jacobi equation:**

$$\begin{aligned}
\int_0^t \left( p \frac{dq}{d\tau} - \mathcal{H} \right) d\tau &= \int_0^t \left( P \frac{dQ}{d\tau} - \mathcal{H}' + \frac{dF(q, P)}{dt} \right) d\tau \\
&= \int_0^t \left( P \frac{dQ}{d\tau} - \mathcal{H}' \right) d\tau \\
&+ F(q'', P'', t) - F(q', P', 0)
\end{aligned} \tag{A.72}$$

$$\frac{\partial(P, Q)}{\partial p, q} = 1 \tag{A.73}$$

consider the transformation for which the new Hamiltonian is zero.  
one gets the Hamilton-Jacobi equation

$$H \left( q, \frac{\partial F}{\partial x}, t \right) + \frac{\partial F}{\partial t} = 0 \tag{A.74}$$

Since  $\mathcal{H}' = 0$ , the new coordinates  $Q$  and  $P$  are constants of motion.

• **Hawking, Stephen:**



Figure A.22: Stephen Hawking.

- **Hawking radiation:**  $T = \kappa$
- **Heisenberg representation:** A Heisenberg state is can be viewed as the value Schroödinger state at some fixed time.
- **Heisenberg uncertainty principle:** From “THE LIFE OF THE COSMOS” by Lee Smolin p.306:

“Let us imagine that we have a friend who lives in Quantumland, where big things can be simple. To tempt us with the beauty of her world, she has sent us a present. We go to the airport to receive it and are given a sealed box with a door on each end. On top of it is written, ‘QUANTUM PET CARRIER - ONLY OPEN ONE END AT A TIME.’ In Quantumland no one would have to be told that you can only open one end of a box at a time, but we will see that the reason to be grateful for this advice.

Bringing the box home, we hurriedly open it to see what is inside. Opening one end, we see the head of a cat peer out! Lovely, but the cat stays inside. It seems that one property of pets in Quantumland is that they can never come out of their box, one can only interact with them by opening one of the doors. OK, we can live with this, but we become curious to at least know the sex of our cat. Well, we can use the door at the other end for this. We try to open it, but we find it is closed tight. Remembering what was written on top, we close the first door. Immediately the back door comes open. By looking in, we are able to ascertain that our pet is a boy.

This done, we go back to the first door to play with our cat. To do this we must first close the back door. We then open the front door to find a jolly looking puppy gazing back at us!

After some trials and examinations, we discover that we are in an interesting situation. When we open the first door we discover that our quantum pet is either a cat or a dog. If we open the second door, we discover that our pet is either male or female. However, by the peculiar properties of the box, our vision is obscured so we cannot be sure, when gazing in the front, what sex our cat or dog is. And when we look in the back door we can ascertain the sex, but we cannot judge reliably whether it is a dog or a cat.

We cannot have both doors open at once, so we can never be sure of both species of our pet and its sex. Ascertaining one destroys knowledge of the other. If all we do is look at the front end, then once we have seen a cat there, we will always see a cat. If we wish, we may at any time close that door and peer in the other side to learn the pet's sex. Whether it is a cat or a dog, we discover that there is a fifty percent probability that it is male and a fifty percent probability that it is female. But, once we have done that, if we go back to the front, we will not necessarily find a cat, as we did before. For once we have ascertained the sex, the species again gets scrambled, and half the time it will be a cat, and half the time it will be a dog.

What we are experiencing is exactly the Heisenberg Uncertainty Principle. It is happening because a complete description of our quantum pet would include its species and its sex. According to classical science, we ought to be able to take the animal out of the box and see what it is. But a quantum pet can never be removed from its box and, for reasons that are perhaps mysterious, we can only ever observe one aspect at a time.

Perhaps the reader thinks I'm being facetious, or teasing. But no, I am describing what we believe is the general situation we are in when we observe any physical system. The Heisenberg Uncertainty Principle limits the information we can have about any system to always exactly half of the information we need to have a complete description. We

always have some choice of which information we would like to have. But, try as we may, we cannot exceed this limit.”

- **holographic principle:**

- **holonomic equivalence:** two loops  $\alpha$  and  $\beta$  are said to be holonomic equivalent if they have the same holonomy with respect to every connection, i.e.

$$H_A(\alpha) = H_A(\beta) \quad \text{for all } A \text{ in the space of connections } \mathcal{A}. \quad (\text{A.75})$$

The set of all holonomic equivalence classes is called the **holonomic loop space** and is denoted  $\mathcal{HL}$ .

- **holonomy algebra:** denoted  $\mathcal{HA}$ .

- **holonomy-flux \*-algebra:** Holonomies of connections lead to well defined operators on  $\mathcal{H}^0$ .

$$h_\alpha(A) = \mathcal{P} \exp \left( \oint_\alpha A_a ds^a \right) \quad (\text{A.76})$$

The connection is smeared in *1-dimensions*. In LQG the classical electric flux  $E_k(S)$  through a surface  $S$  is the integral of the densitised triad  $\tilde{E}_k^a$  over a two surface

, where  $n_a^S$  is the conormal vector with respect to the surface  $S$ .

The conjugate momentum is dual to a 2-form

$$e_{abi} := \frac{1}{2} \eta_{abc} \tilde{E}_i^c$$

geometrically it is natural to smear in 2-dimensions. It again turns out that these 2-dimensionally smeared fields lead to well defined operators on  $\mathcal{H}^0$ .

$$[E_i]_f := \int_S e_{abi} f^i dS^{ab}$$

$$\frac{\delta h_\alpha(A)}{\delta A_i^a(x)} = \int_0^1 \delta^3(x^j - \alpha^j(t)) \frac{d\alpha^i}{dt} (U_\alpha)_0^t(A) \tau_a (U_\alpha)_t^1(A) dt \quad (\text{A.77})$$

$$\frac{\delta E(S)}{\delta E_a^i(x)} = \int_S dudv n_i \delta(x^j - X^j(u, v)) \tau_a, \quad (\text{A.78})$$

By virtue of the canonical commutation relations, the Poisson brackets turn out to be

$$\{E_a(S), h_\alpha(A)\} = \int d^3x \frac{\delta E_a(S)}{\delta E_b^i} \frac{\delta h_\alpha(S)}{\delta A_i^b}. \quad (\text{A.79})$$

From these last three relations we see that if  $S$  and  $\alpha$  have no common points the integration vanishes. The same happens if  $\alpha$  belongs to  $S$  since in this case vectors tangent to the graph are orthogonal to  $n^i$ . We are led to consider loops with a finite number of intersections with the surface. Consider the case of one edge  $e$  intersecting the surface  $S$

$$\begin{aligned} \{[E_S]_f, (h_e)_{mn}\} &= \lim_{\epsilon \rightarrow 0} [(h_{e_1}^\epsilon)_{ml} \{[E_S]_f, (h_{e(\epsilon)})_{lp}\} (h_{e_2}^\epsilon)_{pn}] \\ &= \lim_{\epsilon \rightarrow 0} \left[ (h_{e_1}^\epsilon)_{ml} \left\{ \int_S f_b(y) \tilde{E}^b(y), \int_{e(\epsilon)} A^a(x) \tau_{lp}^a \right\} (h_{e_2}^\epsilon)_{pn} \right] \\ &= \mp 8\pi G \gamma f_a(P) (h_{e_1} \tau_a h_{e_2})_{mn} \end{aligned}$$

$$\{E_a(S), h_\alpha(A)\} = 8\pi G \gamma \tau_a \sum_{e \in \alpha} h_e(A) o(e, S) \quad (\text{A.80})$$

where  $o(e, S)$  is  $+1, -1$  if the orientations of  $e$  and  $S$  agree or disagree, respectively.

Hence one can define the action of  $E(S)$  on a generic cylindrical function by the expression

$$E_a(S)[f] = \{E_a(S), f(h_\alpha(A))\} \quad (\text{A.81})$$

with  $E_a(S)$  acting as a vector field.

The Poisson bracket between the strip functionals vanishes unless the two strips intersect. If they do, the bracket is given by a sum of slightly generalized strip functionals.

If one attempts to smear connections and triads in 3-dimensions the resulting operators fail to be well defined on  $\mathcal{H}^0$ .

The new algebra provides a distributional extension of the old one.

There is uniqueness theorem on this algebra, see section.

• **Holst action:** The Palatini action plus an extra term multiplied by the inverse of the Barbero-Immirzi parameter which does change the classical equations of motion:

$$S_H(e, A) = \frac{1}{2\kappa} \int_{\Sigma \times \mathbb{R}} \left( \frac{1}{2} \epsilon_{IJKL} e^I \wedge e^J \wedge F^{KL} + \frac{1}{\gamma} e^I \wedge e^J \wedge F^{KL} \right)$$

- **hoop group:** the quotient of the loop group with respect to holonomic equivalence.
- **horismos:** The future horismos of a set  $S$  is defined to be  $E^+[S] := J^+[S]/I^+[S]$ .
- **Husain, :**



Figure A.23: Husain.

- **ideal clock limit:**

- **Immirzi-Barbero parameter:** A certain free parameter,  $\gamma$ , in the canonical quantization of gravity. This parameter is not present in the classical theory - In the classical theory, different values of the Immirzi parameters label equivalent classical theories. However, in the quantum theory, representations with different Immirzi parameters are not unitarily equivalent - arises from an ambiguity in the quantization procedure... It can be fixed by the requirement that the quantum gravity computation reproduce the Hawking value for the entropy of a black hole.

A conflicting assertion Alexandrov *et.al.*: [153] [154]. *“It was thought that the anomaly in question is a physical one so that the Immirzi parameter becomes a new fundamental constant. However, in a series of works [153] it was shown that this is not the case because it is actually a consequence of a diffeomorphism anomaly, whereas there is a quantization which preserves all classical symmetries and leads to results independent of the Immirzi parameter.”.*

- **induced metric:** The induced metric the metric  $q$  a hypersurface picks up from the metric  $g$  of the spacetime the hypersurface is embedded. The induced metric of a hypersurface  $\Sigma$  is obtained by restricting the line element to displacements within the hypersurface  $\Sigma$ .

The induced metric is also called the **first fundamental form**.

Some jargon: The induced metric  $q$  is the pullback of the spacetime metric  $g$ , denoted  $\phi^*g$ , where  $\phi$  is the map from the spacetime manifold  $\mathcal{M}$  to the hyperspace manifold  $N$ , i.e.,  $\phi : \mathcal{M} \rightarrow N$ .

- **inertial frames:** The set of all inertial systems is the set that can be derived from a given one by a Lorentz transformations.

- **inextendable curve:**

- **inflation:** An accelerated expansion of the very early universe. Eliminates the need for contrived and highly special initial conditions.

- **inner product for quantum gravity:** In LQG the inner product on spacially diffeomorphism invariant states is known in closed form. One now wishes to have an inner product on the subset of states that satisfy all the constraints. Such a can be written formally in terms of a “projection operator” by means of spin foams - one of the promises of the Master constraint programme is the construction of a well defined physical inner product. - i.e., the physical inner product, ??from which we can calculate physical expectation values??.

- **integrable:**

integrability of of the system in the sense of Liouville:

(1) the number of independent conserved quantities equals the number of degrees of freedom, and that

(2) these conserved quantities are in involution.

are in involution means  $F_i$

$$\{F_i, F_j\} = 0 \tag{A.82}$$

There are standard methods available for solving the system completely (Hamilton-Jacobi theory, action-angle variables, ...).

- **intensive thermodynamical quantities:**

- **interaction picture:** In the Schrödinger picture, time evolution is governed by the states and their equations. In the Heisenberg picture, time evolution is governed by the observables and the equation  $\frac{dA}{dt} = \{B, \mathcal{H}\}$ .

In the interaction picture we divide the time dependence between the states and the observables. This is suitable for systems with a Hamiltonian of the form  $B_H = B_{H0} + B_{H1}$  where  $A_{H0}$  is time independent. The interaction picture has many uses in perturbation theory.

- **interpretations:** Some seek to interpret quantum mechanics in a way which accords more with our intuition derived from classical mechanics.

reduction of the wave-packet of a microscopic system and a specification of its quantum state by a macroscopic instrument

- **interval:** In the general theory of relativity, the square of the interval is given by:

$$ds^2 = g_{ab} dx^a dx^b, \tag{A.83}$$

where  $g_{ab}$  are the components of the metric tensor, and  $dx^a$  is the differential of the coordinate  $x^a$ . Time-like interval  $ds^2 < 0$ ; Space-like interval  $ds^2 > 0$ .

- **inverse loop transformation:**

- **irreducible representation:**

- **Chris Isham:** In a seminal paper with Ashtekar rigorously constructed the quantum configuration space of distributional connections using GNS theory for  $C^*$  algebras. Also later applied topos methods to develop of a theory of quantum gravity.



Figure A.24: Chris Isham.

- **ISO (3) group:**  $ISO(3)$  denotes the Euclidean group in three dimensions.

The translation group is an abelian group.

Obviously

$$\hat{R}\hat{T}\hat{R}^{-1} = \hat{T}' \tag{A.84}$$

consequently the translation group is an invariant abelian subgroup of the Euclidean group, (translation-rotation group).  $SO(3)$  is another subgroup such that the only element they have in common with the translation group is the identity element  $\mathbf{I}$ . Every element of  $ISO(3)$  can be written in a unique way as  $g = RT$ ,  $R \in SO(3)$ ,  $T \in \mathbf{R}^3$  then  $ISO(3)$  is the semi-direct product of  $\mathbf{R}^3$  and  $SO(3)$ ,

$$SO(3) \otimes_S \mathbf{R}^3. \tag{A.85}$$

We introduce translation generators  $P_a$ ,  $a = 1, 2, 3$ , which satisfy

$$[J_a, P_b] = \epsilon_{abc} P^c, \quad [P_a, P_b] = 0, \quad (\text{A.86})$$

- **ISO (2,1) group:**  $ISO(2,1)$  denotes the Poincare group in three dimensions.

- **isolated horizons:**

Isolated horizons are generalizations of the event horizon of stationary black holes to physically more realistic situations. The generalization is in two directions. First, while one needs the entire spacetime history to locate an event horizon, isolated horizons are defined using properties of spacetime *at the horizon*. Second, although the horizon itself is stationary, the outside space-time can contain time-dependent fields and geometry.

Restriction to spacetimes which admit an internal boundary which has certain boundary conditions.

Type **I** the isolated horizon geometry is spherical

Type **II** the isolated horizon geometry is axi-symmetric. This class includes rotating isolated horizons as well as distorted (due to exterior matter), see [136].

Type **III** Most general include arbitrary distortions

Note that the symmetries refer *only* of the horizon geometry. This weak set of boundary conditions give rise to the zeroth and first laws of black hole mechanics, in fact, isolated horizons can be characterized this way.

Black hole entropy [132].

The action principle and the Hamiltonian description is well defined and the resulting phase space has an infinite number of degrees of freedom. The full spacetime metric need not admit any geometric symmetries even in a neighbourhood of the horizon. They quantize this sector of the full phase space using techniques developed for the full theory.

The problem was reduced to counting the number of ways to puncture a two-dimensional sphere, representing the horizon, which gave the horizon area close to a given value.

Isolated Horizons in Numerical Relativity [131].

- **Kaluza-Klein:** The radius of the extra dimension is taken to be fixed and non-dynamical and not subject to Einstein's field equations. As such there are no processes in which gravitational and electromagnetic effects convert into each other. For the theory to be a true unification, gravity and electro-magnetism should be aspects of one phenomenon.

Another objection is that the theory cannot incorporate fermions of the gauge theory.

- **Kauffman, Louis:**

- **kernel:** the set of (eigenstates) solutions of an operator which have zero as their eigenvalue. or

- **Killing horizon:**

$$\xi^a \xi_a = -1 - \frac{2M}{r} \quad (\text{A.87})$$

Thus outside the horizon ( $r > 2M$ )  $\xi^a$  is timelike, at the horizon ( $r = 2M$ )  $\xi^a$  is null.

- **Killing vector field:** Geometry does not change along a Killing vector field. An isometry of the spacetime. A Killing vector field satisfies the Killing equation

$$\nabla_a K_b + \nabla_b K_a = 0. \quad (\text{A.88})$$

- **Kinematic Hilbert space:**

There are several complementary characterisations of the Kinematic Hilbert space  $\mathcal{H} := \mathcal{H}_\omega$  which are useful in different contexts.

1. Positive linear functional characterisation
2.  $C^*$ - algebraic characterisation
3. Projective limit characterisation

- **KMS condition:** Kubo-Martin-Swinger condition for thermal equilibrium.

$$\langle 0, \beta | A(t) B(t) | 0, \beta \rangle = \langle 0, \beta | B(t') A(t + i\beta) | 0, \beta \rangle \quad (\text{A.89})$$

$$\omega((\gamma_t A) B) = \omega(B(\gamma_{t+i\beta} A)) \quad (\text{A.90})$$

Haag *et al* [53] have shown that this KMS condition reduces to the well known Gibbs condition

$$\omega(\cdot) = \frac{\sum_n e^{-\beta E_n} |m \psi_n \rangle \langle \psi_n|}{(\sum_n e^{-\beta E_n} \langle \psi_n | \psi_n \rangle)} \quad (\text{A.91})$$

- **KMS states:** The relevant states describing thermal equilibrium are the so-called KMS-states.

- **Kochen-Specker theorem:** [363] [361] Implies it is not possible to assign actual, *possessed*, values to all quantum mechanical physical quantities of an individual system. (See section contextual interpretations)

It is the one of the no-go theorem against hidden variables theories, (the other being Bell's theorem which states that, given the premise of locality of physical theories, a hidden variables theory cannot match the statistical predictions of quantum mechanics).

better version: [362] Kochen-Specker theorem asserts the impossibility of associating real values  $V(\hat{A})$  to all physical quantities in a quantum theory (if  $\dim\mathcal{H} > 2$ ) whilst preserving the 'FUNC' rule that  $V(f(\hat{A})) = f(V(\hat{A}))$  - i.e., the value of a function  $f$  of a physical quantity  $A$  is equal to the function of the value of the quantity. Equivalently, it is not possible to assign true-false values to all the propositions in a quantum theory in a way that respects the structure of the associated lattice of the projection operators.

Before we can state the theorem we first need to introduce the idea of the valuation of an operator  $\hat{A}$  representing a physical quantity,  $V_s(A)$ . Valuation is assigning truth-values to propositions ' $A \in \Delta$ ' asserting that the quantity  $A$  lies in the Borel subset  $\Delta$  of the spectrum of the operator  $\hat{A}$  that represents  $A$ .

The Kocher-Specker theorem asserts that, in the case when the dimension of Hilbert space is greater than 2, one *cannot* assign real numbers to all quantum quantities in a way that preserves functional relations between them. By 'preserves functional relations' is meant for any operator  $\hat{A}$  and any function of it  $f(\hat{A})$  ( $f$  a function from  $\mathbb{R}$  to  $\mathbb{R}$ ), the value of  $f(\hat{A})$  is the corresponding function of the value of  $\hat{A}$ , i.e.,  $V_s(f(A)) = f(V_s(A))$ .

(1)

??? (2) ideal measurement of an observable  $\hat{A}$  in a state  $\psi \in H$  yields an eigenvalue of  $\hat{A}$  and, immediately after the measurement, the state is thrown into corresponding eigenstate. ???

- **Komar integral:** An expression for the conserved mass and angular momentum for stationary spacetime evaluated when the spacial boundary is pushed to infinity.

$$E = \frac{1}{4\pi G} \int_{\partial\Sigma} d^2x \sqrt{q} n_a \sigma_b \nabla^a K^b, \quad J = -\frac{1}{8\pi G} \int_{\partial\Sigma} d^2x \sqrt{q} n_a \sigma_b \chi^a K^b \quad (\text{A.92})$$

where  $\partial\Sigma$ ,  $\sigma_b$  (timelike and future pointing) and  $n^a$  timelike orthogonal unit normal vectors to  $\partial\Sigma$ . where  $K^b$  time translations for energy.

- **Krasnov, Kirill:**

- **Kruskal coordinates:** These coordinates cover the entire spacetime manifold of a maximally extended solution of classical general relativity.

- **Kuchar's "reduction ad absurdum":** We require a "global time coordinate"

- **the landscape:**

- **lapse function  $N$ :**  $N$ , together with the shift function  $N^a$ , only serves to specify the foliation of  $\mathcal{M}$  and as such it is not itself a dynamical field.




Figure A.25: An expression for the conserved mass, evaluated when the spacial boundary is pushed to infinity. (b)  $\partial\Sigma$  is a closed spacelike two-surface surrounding the source.



Figure A.26: Kirill Krasnov.

- **lattice gauge theories:**

The difference between lattice approaches in Minkowskian field theories and background independent field theories that use floating graphs. An approximation to the continuum theory limit . Whereas in background independent field theories

continuum limit used in lattice approaches in Minkowskian field theories, whereas in background independent field theories allow all possible graphs in  $\mathcal{M}$ .

lattice gauge theory formulations applied to GR conflict with its invariance under diffeomorphisms as it introduces a fixed background geometric structure.

- **Laws of black hole mechanics:**

**The zeroth law of black hole mechanics** stationary black holes have constant surface gravity on the entire horizon.

**The first law of black hole mechanics** tells us how the area of a black hole increases when we make a transition from an initial stationary solution to a nearby stationary

solution.

$$\delta M = \frac{\kappa}{8\pi G} \delta a + \Omega \delta J, \tag{A.93}$$

where  $\kappa$  is the surface gravity.

**The second law of black hole mechanics** states that the area of an event horizon can never decrease.

A system in equilibrium will have settled to a stationary state.

**Isolated horizons:** all quantities which enter the statement of the law now refer to the horizon itself.

- **Laws of thermodynamics:**

**The zeroth law of thermodynamics** states that in thermal equilibrium the temperature is constant throughout the system.

**The first law of thermodynamics** tells us how the

$$\delta E = T \delta S + \delta W. \tag{A.94}$$

**The second law of thermodynamics** states that the entropy of an isolated system can never decrease.

- **Lax pairs:**

$$\frac{dL}{dt} \equiv \dot{L} = [M, L]. \tag{A.95}$$

$$L(t) = g(t)L(0)g(t)^{-1} \tag{A.96}$$

where the invertible matrix is determined by the equation

$$M = \frac{dg}{dt} g^{-1}. \tag{A.97}$$

- **Lewandowski, Jerzy:**

- **Lie derivative:** Lie derivative.

see maths glossary.

- **linear 2-cell spin-foam models:**




Figure A.27: Jerzy Lewandowski.

- **Livine, Etera:**



Figure A.28: Etera Livine.

- **local particles:** Realistic particle detectors are finitely extended. Physical states detected by a localized detector (eigenstates of local operators describing detection) are local particles states. Local particles have a chance of retaining meaning in the fully general relativistic context where Fock states are not available.

- **Loll, Renate:** Loll works on developing a theory of quantum gravity, reconciling the beautiful geometric description of space and time laid out in Einstein's theory of General Relativity with the insight that all of physics at its most fundamental level must be described by quantum laws of motion. She is one of the pioneers of a new approach to the nonperturbative (background independent) quantization of gravity, that of Causal Dynamical Triangulations which recently has produced a number of remarkable results.

These include a dynamical derivation of the fact that space-time is four-dimensional (something that can be taken for granted only in classical gravity) and that it has the shape of a de Sitter Universe (like our own universe in the absence of matter), and of the so-called wave function of the universe which plays an important role in understanding the quantum behaviour of the very early universe. Remarkably, one also finds that the dimensionality of spacetime reduces smoothly to two at short distances, indicative of a highly nonclassical behaviour of spacetime geometry near the Planck scale. These results are obtained by superposing elementary quantum excitations of geometry which have a notion of causality (“cause preceding effect”) built into them at the very smallest scale.



Figure A.29: Renate Loll.

• **loop algebra:**

given a connection the identities satisfied by Wilson loops may be implemented on the free vector space of formal sums of *single* loops by requiring that if

$$\sum_i c_i T[\alpha_i, A] = 0 \tag{A.98}$$

for *all* connections, then these loops are linearly dependent

$$\sum_i c_i \alpha_i = 0 \tag{A.99}$$

on the free vector space of single loops.

When the product on the free vector space is defined by

$$\left(\sum_j c_j \alpha_j\right) \cdot \left(\sum_k d_k \alpha_k\right) = -\sum_{j,k} c_j d_k \left(\alpha_j \# \beta_k + \alpha_j \# \beta_k^{-1}\right) \quad (\text{A.100})$$

these Mandelstam relations define an ideal. The quotient space by the ideal defines an algebra, which is called the holonomy algebra...

[gr-qc/9512020]

- **loop quantum cosmology:** Mini-superspace but which appropriate incorporation of the discrete nature of quantum geometry. Very good accessible account can be found in [241].

As yet it makes no attempt to address the problem of time or interpretation of the wavefunction of the universe.

- **loop quantum gravity LQG:**

A background independent approach to quantum gravity. In many ways conservative based only on the experimentally well confirmed principles of general relativity and quantum mechanics.

a canonical quantization of a Hamiltonian description of classical general relativity, whereby spacetime is sliced into a stack of 3-d spacial hypersurfaces Arnowitt-Deser-Misner (ADM). In LQG the basic variable used is, rather than a metric, a connection and its “conjugate momentum” a triad field (can be thought of as the square root of the 3-d spatial metric). The spacial 3-metric on a hypersurface is constructed from the triad field and the connection contains information about extrinsic curvature of the slice of the slice sitting in the spacetime, establishing contact with the metric ADM formulation.

Wilson loops, polymer nature

- **loop representation:** import machinery from particle physics... [54].

- **loop transformation:** the Fourier transformation [54]

$$\tilde{\psi}(k) = \int dk e^{ik \cdot x} \psi(x), \quad (\text{A.101})$$

which relates the position representation with the momentum representation.

$$\psi(\gamma) := \int_{\mathcal{A}/\mathcal{G}} d\mu(A) T^0(\gamma)[A] \Psi(A), \quad (\text{A.102})$$

where the Wilson loop functionals  $T^0$  plays the role of an integral kernel, (as does  $e^{ik \cdot x}$  in the Fourier transformation), and wave functions  $\psi(\gamma)$  in the loop representation are

labelled by closed spacial curves  $\gamma$ . Once the loop transform is rigourously defined the loop representation is unitarily equivalent to the connection representation.

- **Lorentz covariant loop quantum gravity (CLQG):** Standard loop quantization is based on the Ashtekar-Barbero formulism and leads to the theory in which  $SU(2)$  gauge group in the tangent space. It starts with the first order formulation of general relativity in 3+1 dimensions with the Lorentz gauge group in the tangent space. Then as a result of partial gauge fixing, the canonical formulation possesses, besides the usual diffeomorphism invariance, only a local  $SU(2)$  symmetry.

- **LOST uniqueness theorem:** Named after the people who developed it: Lewandowski, Okolow, Sahlmann, and Thiemann. In LQG once one decides to base quantization on the holonomy-flux algebra and have the spacial diffeomorphism group implemented unitarily, there is only one choice for the Kinematical Hilbert space. Once the Kinematical representation has been chosen in which the constraints can be defined as operators, the physical representation follows by a rather tight procedure. Hence, the whole quantization programme is put on a very theoretical robust footing leading to a falsifiable theory of quantum gravity. One prediction of the theory being the discreteness of the eigenvalues of geometric operators.

- **luminosity distance:**

- **LSZ reduction formulas:** using the LSZ reduction formulas it is more easily demonstrated that the a scattering amplitude is unchanged by making in states  $|in\rangle$  into out states  $|out\rangle$  and vice versa. so we have a symmetry that implies we can exchange preparation of states and the measurement proper.

- **Mach's principle:**

- **macroscopics:**

- **Major, Seth:**

- **marginally trapped surfaces:** See trapped surface.

- **Markopoulou, Fontini:**

- **Master constraint:** The multiplicity of smeared Hamiltonian constraints combined into one constraint.

since only a non-anomalous constraint algebra usually leads to a sufficiently large enough physical Hilbert space  $\mathcal{H}_{Phys}$

the question of anomaly freeness is translated into the size of the physical Hilbert space  $\mathcal{H}_{Phys}$ .

the infinite dimensional constraint algebra, whether it closes with structure functions or structure constants, is replaced by a one-dimensional Abelian Lie algebra.



Figure A.30: Fontini Markopoulou.

- **Master equation:** Any complete observable  $\mathcal{O}$  must be constant under a transformation generated by the Hamilton constraint  $C_j(x)$  i.e.,  $\{\mathcal{O}, C_j(x)\}|_{C=0} = 0$ . This condition is equivalent to

$$\{\mathcal{O}, \{\mathcal{O}, \mathbf{M}\}\}_{\mathbf{M}=0} = 0 \iff \{\mathcal{O}, C_j(x)\}|_{C=0} = 0. \quad (\text{A.103})$$

see Eq (??).

- **Master Constraint Operator:** Replaces the Hamiltonian constraints  $\hat{C}_a$  by a single Master constraint  $\hat{\mathbf{M}}$  which is a spacially diffeomorphism invariant integral of squares of the individual Hamiltonian constraints

$$\hat{\mathbf{M}} := \frac{1}{2} \hat{C}_a Q^{ab} \hat{C}_b \quad (\text{A.104})$$

which encodes all the necessary information about the constraint surface and the associated invariants (Dirac observables).

- **Master Constraint Programme:**

**Extended Master Constraint Programme:**

- **maximal extension of a spacetime:** in the context of the Schwarzschild spacetime coordinate systems that allow the continuation of a metric across the event horizon. There also is an inner horizon. is maximally extended if the coordinate system allow for continuation of the metric across every horizon.??

- **matrix models:** field theories over groups.
- **maximally extended curve:** A timelike curve is maximally extended in the past if it has no past endpoint. (Such a curve is also called past-inextendible). The idea behind this is that such a curve is fully extended in the past direction, and not merely a segment of some other curve.
- **mesoscopics:** gravity experiments quantum mesoscopics and structural foundations of quantum mechanics.
- **microscopics:**
- **moments:**

$$\frac{1}{|\mathbf{x} - \mathbf{x}'|} = \frac{1}{r} + \frac{\mathbf{x} \cdot \mathbf{x}'}{r^3} + \frac{1}{2r^5}(3(\mathbf{x} \cdot \mathbf{x}') - r^2 r'^2) + \dots \quad (\text{A.105})$$

With this expansion, we can rewrite  $\Phi(\mathbf{x})$  as

$$Q_{ij} = \int d^3x' \rho(\mathbf{x}') (3x'_i x'^j - r'^2 \delta_{ij})$$

- **Montevideo interpretation:** In this interpretation environmental decoherence is supplemented with loss of coherence due to the use of realistic clocks to measure time to solve the measurement problem. The resulting formulation is framed entirely in terms of quantum objects without having to invoke the existence of measurable classical quantities like time in ordinary quantum mechanics. The formulation eliminates any privileged role to the measurement process giving an objective definition of when an event occurs in a system.

- **M-theory:** background independent M-theory for strings

does not presume that they are vibrating in an preexisting spacetime. a formulation in the absence of spacetime in which instead spacetime emerges from the collective behaviour of the strings.

- **M-theory conjecture:** There is a background independent formulation of string theory which unifies all the known string theories, 11 dimensional supergravity and the 11 dimensional supermembrane theory.

C. M. Hull and P.K. Townsend, Unity of superstring dualities, Nucl. Phys. B348 (1995) 109; P. Townsend, (M)embrane theory on T 9, Nucl. Phys. (Proc. Suppl) 68 (1998) 11-16; hep-th/9507048, in Particles, Strings and Cosmology, ed J. Bagger et al (World Scientific, 1996); hep-th/9612121; I. Bars, hep-th/9608061; hep-th/9607122; Petr Horava, MTheory as a Holographic Field theory, hep-th/9712130, Phys.Rev.D59 (1999)046004.

E. Witten, String theory in various dimensions, Nucl. Phys. B443 (1995) 85.

Once (or if) one finds this theory, one must find the classical solutions to this theory and show that the different perturbative string theories arise from expansions around them.

- **multipoles:**

**field multipoles**

**source multipoles**

- **naked singularities:** . In recent paper [398] Harada ...*We propose the concept of ‘effective naked singularity’, which will be quite helpful because general relativity has the limitation of its application for the high-energy end. The appearance of naked singularities is not detestable but can open window for new physics of strongly curved spacetime.* - which of course includes quantum gravity.

- **negative mass:** (the so-called Benny Hill effect).

- **noiseless subsystems:** The framework of noiseless subsystems has been developed as a tool to preserve the fragile quantum information against decoherence []. In brief, when a quantum register (a Hilbert space) is subjected to decoherence due to an interaction with an external and uncontrollable environment, information stored in the register is, in general, degraded. It has been shown that when the source of decoherence exhibits some symmetries, certain subsystems of the quantum register are unaffected by the interactions with the environment and are thus noiseless. These noiseless subsystems are therefore very natural and robust tools that can be used for processing quantum information.

- **net of operator algebras:** an open region of spacetime  $\omega \subset \mathcal{M}$

$$\omega \subset \omega' \implies \mathcal{A}_\omega \subset \mathcal{A}_{\omega'} \quad \text{for all } \omega, \omega' \subset \mathcal{M}, \mathcal{A}_\omega, \mathcal{A}_{\omega'} \subset \mathcal{A}, \quad (\text{A.106})$$

- **Newmann-Penrose null tetrads:** frames that consisted of two real null vectors and two “complex null vectors”. seemingly strange choice well suited for analysis of null geodesics. Intimately related to spinor basis

- **Noether’s theorem:** If the equations of motion possesses some continuous symmetry, then there will be a conservation law associated with that symmetry. The most familiar examples of conservation laws are:

- (i) if there is invariance under some *spatial translation*, then *momentum* is conserved;
- (ii) if there is invariance under *time translation*, then *energy* of the system is a conserved quantity;
- (iii) if there is invariance under *rotation* about some axis, then *angular momentum* about the same axis is conserved.

Noether's gave a proof of the general result. Most physical theories implement dynamics as a result of a variation principle, by means of a Lagrangian. Continuous transformations that leave the action invariant - except for boundary terms.

**(i) Spatial translation:** using the Lagrangian  $\mathcal{L}(x, t)$ .

$$\mathcal{L}(x + \delta x, t) - \mathcal{L}(x, t) = 0 \Rightarrow \frac{\partial \mathcal{L}}{\partial x}(x, t) = 0 \quad (\text{A.107})$$

using this in the Euler-Lagrange equations (E.7),

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{x}}(x, t) = 0 \Rightarrow \frac{\partial \mathcal{L}}{\partial \dot{x}}(x, t) =: p = \text{Const}' \quad (\text{A.108})$$

**(ii) Time translation:** That the Lagrangian does not explicitly depend on  $t$ ,

$$\frac{\partial \mathcal{L}}{\partial t}(x, t) = 0 \quad (\text{A.109})$$

$$\frac{d\mathcal{L}}{dt} = \frac{\partial \mathcal{L}}{\partial t} + \dot{x} \frac{\partial \mathcal{L}}{\partial x} + \ddot{x} \frac{\partial \mathcal{L}}{\partial \dot{x}} \quad (\text{A.110})$$

Making use of the Euler-Lagrange equations

$$\frac{d\mathcal{L}}{dt} = \dot{x} \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{x}} + \ddot{x} \frac{\partial \mathcal{L}}{\partial \dot{x}} = \frac{d}{dt} \left( \dot{x} \frac{\partial \mathcal{L}}{\partial \dot{x}} \right), \quad (\text{A.111})$$

$$\frac{d}{dt} \left( \dot{x} \frac{\partial \mathcal{L}}{\partial \dot{x}} - \mathcal{L} \right) = 0. \quad (\text{A.112})$$

$$E = \dot{x} \frac{\partial \mathcal{L}}{\partial \dot{x}}(x, \dot{x}) - \mathcal{L}(x, \dot{x}) \quad (\text{A.113})$$

• **non-commutative geometry:** The idea is to describe geometry in algebraic terms. One deals with a function algebra over the space one is interested in and the geometry of a manifold is reformulated in terms of this algebra. A topological space  $\mathcal{M}$  is completely described by the commutative algebra  $\mathcal{C}^0(\mathcal{M})$  of continuous functions over it. Once the algebraic description of the geometry is obtained, the structure algebra can be made noncommutative.

Noncommutative geometry is the study of noncommutative algebras as if they were algebras of functions on spaces. noncommutative algebras should be studied as if they were the function algebras of "noncommutative spaces"

See for example [401] for introduction to mathematical constructions.

Connes proved that there is an equivalence between a spinorial compact manifold and a Spectral Triple  $(\mathcal{C}(\mathcal{M}), \mathcal{D}, \Gamma)$ , which contains the algebra, the Dirac operator  $\mathcal{D}$  encoding the differential structure and the metric, and finally the chiral operator  $\Gamma$ .

Connes. quantum mechanics does not permit the measurement of distance smaller than the Plank length and time intervals smaller than the Plank time. Non-commutative spacetimes are adopted as *background* spacetimes.

- **non-commutative spacetime:** non-commutative spacetime can be derived from a quantum deformation which replaces the enveloping algebra of the Poincare Lie algebra by a non-commutative Hopf algebra(quantum group).

□

- **non-commutative theories:** hep-th/9808192

“One success of noncommutative geometry is that the standard model of particle physics of particle physics. As Alain and his colleagues discovered, if you take Maxwell’s theory of electrodynamics and write down the simplest possible noncommutative geometry, out pops the Weinberg-Salam model unifying electromagnetism with the weak nuclear force. In other words, the weak interactions, together with the Higgs field, show up automatically and correctly.”

Non-commutative geometry is a third approach to quantum gravity. It is by definition a background independent approach, as the basic idea is to replace the background manifold by algebraic generalisations of a certain set of diffeomorphism invariant observables, which are the spectrum of the Dirac operator. □

- **non-expanding horizon (NEH):** A horizon is a compact, spacelike 2-surface (usually a sphere) expanding at the speed of light, however, not changing its area element.

The zeroth and first law of Black Hole thermodynamics still hold

A three dimensional sub-manifold  $\Delta$  of spacetime is said to be non-expanding horizon (NEH) if it satisfies the following conditions:

- (i)  $\Delta$  is topologically  $S^2 \times \mathcal{I}$  and null where  $\mathcal{I}$  is an interval on the real line;
- (ii) The expansion  $\theta_{(\ell)} := q^{ab} \nabla_a \ell_b$  of  $\ell$  vanishes on  $\Delta$ , where  $\ell$  is any null normal to  $\Delta$  and  $q_{ab}$  is the degenerate metric on  $\Delta$ ;
- (iii) All equations of motion hold at  $\Delta$ .

- **non-perturbative:** does not describe dynamics of linear perturbations around some fixed background spacetime.

compare with background independence: does not distinguish any particular background metric from the outset.

- **non-renormalizability:**

Part of the reason why some people consider non-renormalizability unacceptable as a physical theory may lie in what one considers the meaning of “quantization” to be.

- **null-tetrads:** Newman-Penrose null-tetrads.

$$\ell^a := \frac{1}{\sqrt{2}}(r^a + t^a), \quad n^a := \frac{1}{\sqrt{2}}(r^a - t^a) \quad (\text{A.114})$$

$$\ell^2 = 0, \quad n^2 = 0, \quad (\text{A.115})$$

complex null vectors

$$m := \frac{1}{\sqrt{2}}(e_1 + ie_2), \quad \bar{m} := \frac{1}{\sqrt{2}}(e_1 - ie_2) \quad (\text{A.116})$$

- **normal region:** A normal region

- **observables:** Observables in Hamiltonian mechanics are real functions defined on phase space,  $\mathbb{R}^{2n}$ . Some examples of observables are position, momentum, energy, angular momentum.

Observables is a property of a physical system that can in principle be measured. In quantum mechanics, an observable is represented by a *self-adjoint operator*.

In quantum mechanics, however, an observable is represented by a self-adjoint operator defined on a dense subspace of a Hilbert space. Work out mean values, mean deviations - the operator to any power should be a finite quantity for it to have a physical meaning.

In the classical GR we compare observations with coordinate-independent quantities. The coordinate time (as well as the spatial coordinates) can in principle be discarded from the formulation of the theory without loss of physical content, because results of real gravitational experiments are always expressed in coordinate-free form.

- **Oeckl, Robert:** In particular, he works on an extension of the standard formulation of quantum mechanics, precisely to make it compatible with general relativistic concepts. This program is called the general boundary formulation of quantum theory.

- **off-shell:**

- **one-partial irreducible (1PI):** See Feynmann diagrams.

- **on-shell:** Equations of motion are imposed.

**off-shell**



Figure A.31: Robert Oeckl.

- **operator:** An operator is a linear map taking vectors to vectors

$$\mathbf{A} : |\psi\rangle \rightarrow \mathbf{A}|\psi\rangle$$

linear meaning

$$\mathbf{A}(a|\psi\rangle + b|\phi\rangle) = a\mathbf{A}|\psi\rangle + b\mathbf{A}|\phi\rangle$$

- **operator algebra:**

$$[\hat{q}, \hat{p}] = i. \tag{A.117}$$

Define

$$U = 1 + i\delta q' \hat{p}, \tag{A.118}$$

where  $\delta q'$  is an infinitesimal parameter. To infinitesimal order this is a unitary operator,

$$U^\dagger U = (1 - i\delta q' \hat{p})(1 + i\delta q' \hat{p}) = 1 + \mathcal{O}((\delta q')^2), \tag{A.119}$$

Hence  $U$  preserves the norm of operators it acts on. It is easily seen that  $U$  is the operator which performs infinitesimal translations on the coordinate operator  $\hat{q}$ :

$$UqU^{-1} = q + i[p, q]\delta q' = q + \delta q', \quad (\text{A.120})$$

therefore

$$\langle q'|Uq = \langle q'|UqU^{-1}U = \langle q'|(q + \delta q')U = (q + \delta q') \langle q'|U, \quad (\text{A.121})$$

which implies that

$$\langle q'|U = \langle q + \delta q'|. \quad (\text{A.122})$$

• **operator ordering ambiguity:** Hamiltonian  $\mathcal{H}(q, p) = p^2q$ . The classical variables replaced by their operators  $p \rightarrow \hat{p} = -i\hbar\frac{\partial}{\partial q}$  and  $[\hat{p}, \hat{q}] = i\hbar$ . There are two possible hermitian Hamiltonian operators corresponding to the classical Hamiltonian,

$$\begin{aligned} \hat{\mathcal{H}} &= \frac{1}{2}(\hat{p}^2\hat{q} + \hat{q}^2\hat{p}) \quad \text{and} \quad \hat{\mathcal{H}}' = \hat{p}\hat{q}\hat{p}, \\ \hat{\mathcal{H}} &= \hat{\mathcal{H}}' - i\hbar\hat{p}. \end{aligned}$$

Operator ordering ambiguities is an important issue in LQG as it relates to the size of the physical Hilbert space, which in turn relates to the question of whether LQG has the correct semi-classical limit.

• **operator valued distribution:**

$$\hat{A}^a(x) = \int_{-\infty}^{\infty} dk e^{ix \cdot k} \hat{a}_k \quad (\text{A.123})$$

A quantum field  $\Phi(x)$  at a point cannot be a proper observable. Physically this appears evident because a measurement at a point would necessitate infinite energy. It may be regarded as a sesquilinear form on a dense domain  $\mathcal{D} \subset \mathcal{H}$ ; this means that the matrix element  $\langle \Psi_2 | \Phi(x) | \Psi_1 \rangle$  is a finite number when both  $\Psi_1$  and  $\Psi_2$  are in  $\mathcal{D} \subset \mathcal{H}$  characterized by the property that the probability amplitudes for particle configurations decrease fast with increasing momenta and increasing particle number.

and that it depends linearly on  $\Psi_1$ , conjugate linearly on  $\Psi_2$ . To obtain an operator defined on the vectors in  $\mathcal{D}$  one has to average (“smear out”)  $\Phi$  with a smooth function  $f$  in Minkowski space i.e. take

$$\Phi(f) = \int \Phi(x) f(x) d^4x.$$

- **Osterwalder-Schrader Reconstruction:** An integral of the form “ $\int \mathcal{D}\phi e^{iS[\phi]/d}$ ” oscillates wildly. We instead first consider the integration “ $\int \mathcal{D}\phi e^{iS[\phi]/d}$ ”; If the Lagrangian is positive then the integral will fall off rapidly. If one can define this and it satisfies certain conditions one can go back to the Lorentzian theory with the Osterwalder-Schrader reconstruction.

, most importantly, requires that the measure  $d\mu(\phi)$  be invariant with respect to Euclidean transformations and that

reflection with respect to the hyperplane

$$(x^0 = 0, x^1, \dots, x^d). \tag{A.124}$$

The Osterwalder-Schrader Reconstruction theorem allows one to obtain the Hilbert space of the canonical quantum field theory from the Euclidean one.

The Osterwalder-Schrader reconstruction of Euclidean QFT was generalized to diffeomorphism invariant theories. Replacing the Euclidean group with the diffeomorphism group. [250]

M. Rainer, *Is Loop Quantum Gravity a QFT?*, [gr-qc/9912011].

- **outer marginally trapped surface** See trapped surface.

- **Page-Wooters construction:** Proposed by Page-Wooters [47] to deal with an aspect of the ‘problem of time’. It consists in promoting all variables of the system to quantum operators and choosing one of the variables to be a “clock” and computing conditional probabilities for the other variables to take certain values when the “clock” takes a certain value. This proposal runs into technical difficulties when applied in detail to constrained systems, as emphasized by Kuchar [41]. In constrained systems the physically observable variables are those that have vanishing Poisson brackets with the constraints. If one of the constraints is the Hamiltonian then this candidate for a clock does not evolve. Page-Wooters tried circumvent this difficulty by considering “kinematical” variables, i.e. variables that don’t commute with the constraints and hence appear to “evolve”. This brings about other problems. Such variables can be promoted to quantum operators acting on the space of wavefunctions that are not necessarily annihilated by the constraints (the kinematic Hilbert space). When the zero eigenvalue of a constraint operator lies in the continuous spectrum (as is the case with GR), the states that are annihilated by the constraints are distributional. Therefore it is difficult to see how they can admit a probabilistic interpretation.

- **Palentini first order formulism of GR:**

This condition, which is imposed *a priori* in Einstein’s formulation of general relativity, is seen to be part of the equations of motion.

$$\mathcal{D}e_a = \partial_b e_a + \epsilon_{abc} A^b e^c = 0. \quad (\text{A.125})$$

• **partial Cauchy surface:** A partial Cauchy surface,  $\Sigma$ , for a spacetime  $\mathcal{M}$  is a hypersurface which no causal curve intersects more than once. An important example is that of the hypersurface  $\Sigma$  extending from a black hole horizon to spacial infinity.

• **partial observables:** These are measurable quantities not determined by the theory, however, correlations between them are. These correlations constitute the dynamics of the theory. We understand dynamics as the relative evolution of partial variables with respect to one another - in which all these variables are treated on an equal footing.

By choosing one of these variables to be an *independent* (“time”) and the rest to be *dependent* variables.

e.g. area as described in chapter 1.

doesn't commute with constraints Kuchar.

see complete observables.

• **partial truths:** cosmology. When it will be in the causal past of the point  $p$ , once it is it remains in the past, there are some points that will never be in the past of  $p$ .

• **particle:** Fock states: with no interactions present we can expand  $\hat{\psi}(x)$  as a complete set of normal modes

$$\{u_i(x)a_i + v_i(x)b_i^\dagger\} \quad (\text{A.126})$$

particles are objects that are trigger readings of detector or that produce bubbles in bubble chambers.

• **particle horizon:** Two different regions of the universe that are separated from each other by a distance greater than  $2R_H(t)$  at epoch  $t$  can not be causally related. This boundary is called the “particle horizon”.

• **partition function:** The partition function is

$$Z(\beta, \mu) = \text{Tr} e^{-\beta(\hat{\mathcal{H}} - \mu\hat{N})} \quad (\text{A.127})$$

where  $\hat{\mathcal{H}}$  is the many-body Hamiltonian,  $\hat{N}$  the particle number operator, and  $\mu$  the chemical potential.

$$\mathcal{E} = -\frac{1}{V} \frac{\partial \log Z}{\partial \beta} = \frac{\partial \beta \mathcal{F}}{\partial \beta}, \quad p = \frac{1}{\beta} \frac{\partial \log Z}{\partial V} = -\mathcal{F} \quad (\text{A.128})$$

- **passive diffeomorphism transformation:** Refers to a change of coordinates, i.e. the same object is represented in a different coordinate system.
- **peeling theorem:** asymptotic expansions.
- **Penrose abstract tensor notation:**
- **Penrose inequality:** The Penrose inequality relates the area of cross-sections of an event horizon  $A_E$  on a Cauchy surface with the ADM mass  $M_{ADM}$  at infinity:

$$\sqrt{A_E/16\pi} \leq M_{ADM}. \quad (\text{A.129})$$

The Penrose inequality has been one of major open questions in classical general relativity, closely tied to an even bigger open question, the cosmic censorship conjecture. It was attempts at finding violations of this that led Penrose to formulate this inequality. The derivation relies so heavily on the cosmic censorship conjecture that a violation of this inequality would go a long way towards contradicting it.

- **Penrose-Carter diagram:** Let  $\tilde{\mathcal{M}}$  denote physical space-time with metric  $\tilde{g}_{ab}$ . The idea was to construct another “unphysical” manifold  $\mathcal{M}$  with boundary  $\zeta$  and metric  $g_{ab}$ , such that  $\tilde{\mathcal{M}}$  is conformal to the interior of  $\mathcal{M}$  with  $g_{ab} = \Omega^2 \tilde{g}_{ab}$ , and so that the “infinity” of  $\tilde{\mathcal{M}}$  is represented by the finite hypersurface  $\zeta$ . We realise the whole physical spacetime  $\tilde{\mathcal{M}}$  as a subset of the larger spacetime  $\mathcal{M}$ .

Asymptotic properties of  $\tilde{\mathcal{M}}$  and of fields in  $\tilde{\mathcal{M}}$  can be investigated by studying  $\zeta$ , and the local behaviour of the fields at  $\zeta$  provided the relevant information is conformally invariant.

- **Penrose process:** extraction of energy from a rotating black hole.




Figure A.32: Penrose process.

- **Sir Roger Penrose:** modern techniques in GR e.g. use of spinors plus elegant topological methods. Invented twistor theory as a theory of quantum gravity. Ingredients spinor networks which were later found an very important place in LQG by Smolin and Rovelli.



Figure A.33: Sir Roger Penrose.

- **perennial:** Perennials are classical variables that are observables, that is, have vanishing Poisson brackets with all the constraints.

One of the constraints, the Hamiltonian constraint, generates you can not expect any of them to work as a clock??

- **Petrov classification:** number of principle spinors
- **perturbative quantum gravity:**

*“However, even if it is accepted that we have finite amplitudes for each fixed topology, we are far from finished. The expressions then have to be summed up. Now there is a problem that the sum apparently actually diverges. The intended finite theory is actually not finite after all! This particular divergence does not seem to worry the string theorists, however because they take this series as an improper realization of the total amplitude. This amplitude is taken to be some analytic quantity, with the power series attempting to find an expression for it by ‘expanding around the wrong point’, i.e. about some point which is singular for the amplitude ....????? This could be OK, although the divergence encountered here has been shown to be of a rather uncontrollable kind (‘not Borel-summable’)..... Moreover, if string-theoretic (perturbational) calculations are actually expansions ‘about the wrong point’, then it is unclear what trust we may place in all these perturbative calculations in any case! Thus we do not yet know whether or not string QFT is actually finite, let alone whether string theory, with all its undoubted attractions, really provides us with a quantum theory of gravity.*

The amplitudes they are calculating are not invariant under active diffeomorphisms and hence cannot be an observables of the theory. So can’t have any meaning beyond the semiclassical limit. Cannot be applied to extreme situations that we would like to access

in quantum gravity.

- **phenomenology of quantum gravity:** Gamma-ray bursts,
- **physical coherent states:** Physical coherent states are needed in order to test the classical limit of gauge-invariant observables.
- **Plebanski action:** Used in a construction of spinfoam models.
- **point holonomies:** Scalar fields e.g. Higg’s particle, quantum cosmology transforms as a scalar so not well defined on an edge - instead we consider the exponential at a point

$$\exp(i\mu A^\varphi(x)) \tag{A.130}$$

• **point-identification:** If one manifold is a replica of another we need a point-identification map which relates points in the two manifolds which are to be regarded as the “same”. The correspondence of the points of these two spacetime manifolds is arbitrary and so is a “gauge freedom”.

- **Poisson bracket:**

$$\{A, B\} = \frac{\partial A}{\partial q} \frac{\partial B}{\partial p} - \frac{\partial A}{\partial p} \frac{\partial B}{\partial q} \tag{A.131}$$

• **polymer particle representation:** [241] If one drops the requirement of weak continuity we can circumvent the Stone-von Neumann theorem, one finds that there are unitarily inequivalent representation of CCR algebra (Weyl algebra).

The standard Schrödinger representation representation of the Weyl algebra plays the role of the Fock representation of low energy quantum field theories and the new, unitarily inequivalent representation -called the polymer particle representation- in which states are mathematically analogous to the polymer-like excitations of quantum egometry.

It mimics the loop representation - e.g. it has discrete - momentum not a well defined operator in this representation.

Appears in the mathematics of loop quantum cosmology.

- **Ponzano-Regge:** A partion function for 3d GR.

$$W(\mathbf{j}) = \left\{ \begin{array}{ccc} j_1 & j_2 & j_3 \\ j_4 & j_5 & j_6 \end{array} \right\} \tag{A.132}$$

Witten’s *ISO(3)* Chern-Simons theory [E. Witten, Nucl. Phys. **B311**, 46 (1988).]


$$Z_{PR} = \sum_j \prod_f \dim(j_f) \prod_v$$


Figure A.34: PonRegg.

- **pre-geometry:** More fundamental substratum from which our familiar continuum physics is assumed to emerge as a coarse grained and secondary model theory. The dynamics of pre-geometry in the appropriate physical regime, give rise to the appearance of a background spacetime with matter fields or particles whose dynamics are played out over it. Examples of pre-geometric theories, spinfoam of LQG,...

- **preons:** Quarks, leptons and heavy vector bosons are suggested to be composed of a stable spin-1/2 preons, existing in three flavours, combined according to simple rules.

- **prequantization:** A prequantisation of a symplectic manifold  $(M, \omega)$  is a pair  $(L, D)$  where  $L$  is complex Hermitian line bundle over  $M$  and  $D$  a compatible connection with curvature  $(1/\hbar)\omega$ .

- **primary constraints:**

- **principal fibre bundle:** Loosely speaking, in the context of gauge theory in spacetime, it is a manifold that locally looks like cartesian product of spacetime manifold and the parameter space of the gauge group. A fibre bundle whose fibre is a Lie group in a certain special way.

Principal bundles are equivalent, if the fibres are transform  $g \rightarrow hg$ . This corresponds to a gauge transformation for the connection 1-form,  $A_\mu(x)$ . The curvature of this bundle connection then turns out to be the Maxwell field tensor  $F_{ab}$ .

- **principle of complementarity:** The predictions of general relativity will coincide with the Newtonian gravity when there particles have small mass and speeds are small compared to the speed of light  $c$ .

- **problem of time:** The problem of time in quantum gravity is a bit tricky to describe, since it takes different guises in different approaches to quantum gravity. (beaz week 41)

In a time reparameterization invariant system of General Relativity there is no time evolution - “nothing happens” in the theory in obvious contradiction to what we observe.

Newton’s absolute time is an apparently indispensable notion in standard quantum mechanics. But GR teaches us that there is no external time - no true time by which mechanics evolves. There is only relative evolution of variables. Time must be promoted to the same status with other variables.

In the first chapter we saw that reference bodies are necessarily part of this dynamical system under study - so system is closed and all dynamical variables behaves quantum mechanically - all variables at the same level. In the context of closed systems where everything behaves quantum mechanically. also in quantum cosmology close to the big bang.

- **projected spin network:**

- **proper length:** a proper length of an object is its length as measured in the rest system of the object?????????

- **proper time:**The label of “the problem of time” is often given to a number of related, but slightly different issues.

Briefly put, the problem of time is as follows: how is one to apply quantum mechanics to general relativity in which a classical non-dynamical background time is missing?

the proper time of an object is the time measured by a clock located in the rest system of the object.

- **Pullin, Jorge**



Figure A.35: Jorge Pullin.

- **qef representation (quantized electric Flux representation):** Analogue of spin network states for background independent Maxwell theory. Electric field flux is quantized. The analogous result in  $SU(2)$  LQG theory , with its geometric interpretation, corresponds to quantized area.

- **quadrupole moment:** Einstein already in 1917 worked out for quadrupole

$$Q_{ij} = \int d^3x' \rho(\mathbf{x}') (3x'_i x'^j - r'^2 \delta_{ij})$$

See moments.

- **quantization:** [arXiv: quant-ph/ 0412015]. Quantisation is the problem of deriving the mathematical framework of a quantum mechanical system from the mathematical framework of the corresponding classical mechanical system. A method of quantisation must contain a map  $\mathcal{A}$  from the set of classical observables to the set of quantum observables with the following properties:

The theories exist in their own right and perturbation methods serve as approximation techniques to extract answer to "physically interesting questions."? hep-th/9408108

- **quantum causal histories:** [367]

???Sorkin. Postulating that the fundamental underlying structure is a discrete causal. Attempts to recover the continuum picture by suitable coarse graining.???

- **quantum configuration space:** In quantum mechanics systems with a finite number of degrees of freedom, states are represented by functions on the classical configuration space. However, in field theories quantum states are on a larger space - the *quantum configuration space*.

For systems with a finite number of degrees of freedom, the classical configuration space also serves as the space of independent variables of the wavefunction of the quantum theory; we say that the configuration space is then the quantum configuration space.

For systems with an infinite number of degrees of freedom, on the other hand, the quantum configuration space is an enlargement of the classical configuration space. The free field theory in Minkowski spacetime while the classical configuration space can be built from suitably continuous fields, the quantum configuration space includes all tempered distributions.

$$\int [\mathcal{D}\varphi] e^{iS} \Phi(x) \Phi(x) \tag{A.133}$$

Typical measures used in quantum field theories are. The classical configuration spaces are contained in sets of measure zero. As a result, the action  $S$  inside the functional integral is ill-defined

In the Hilbert space language, this is the origin of field theoretic infinities, e.g., the reason why we can not naively multiply field operators.

The presence of an infinite number of degrees of freedom causes only one major modification: the classical configuration space  $\mathcal{C}$  of smooth fields is enlarged to the quantum

configuration space  $\mathcal{S}'$  of (tempered) distributions. Quantum theoretical difficulties associated with defining products of operators can be directly traced back to this enlargement [34].

This enlargement from  $\mathcal{A}$  to  $\overline{\mathcal{A}}$  which occurs in the passage to the quantum theory is very similar to the enlargement from  $\mathcal{C}$  to  $\mathcal{S}'$  in the case of scalar fields. This enlargement plays a key role in the quantum theory (especially in the discussion of surface states of the quantum horizon).

- **quantum cosmology:** A theory that attempts to describe the whole universe in quantum-mechanical terms.

- **quantum field theory: Background *dependent* quantum field theory:** In spite its many practical successes where predictions from calculations have been shown to agree with experiment with phenomenal accuracy, QFT is mathematically a problematic construction.

very naturally defined, that we are however unable to calculate without incurring infinities.

can be reformulated in terms of classical statistical field theory.

**Background *independent* quantum field theory:**

- **quantum field theory on curved spacetime:** There is often confusion between a dynamical theory *on* a given curved spacetime with the dynamic theory *of* the spacetime, which is what GR is really about.

From [304]:

“Approximation methods for gauge invariant observables provide an explicit way to construct observables and may also help to test (quantum) interpretation of these as well as to discuss phenomenological implications [8, 9]. The approximation scheme developed in this work shows explicitly how to express local observables in a gauge invariant way and therefore indicates how one could understand quantum field theory on curved space time as an approximation to the full theory of quantum gravity.”

- **quantum gauge fixing:** An approximation scheme

all the degrees of freedom are fluctuating but the unphysical ones’ fluctuations are suppressed by the semiclassical state.

- **quantum mechanics:**

- **quantum logic reconstruction:** Quantum logic reconstruction attempts

From the single premise that the “experimental propositions” associated with a physical system are encoded by projections in the way indicated above, one can reconstruct the

rest of the formal apparatus of quantum mechanics. The first step, of course, is Gleason's theorem, which tells us that probability measures on  $L(H)$  correspond to density operators.

Taken directly from <http://plato.stanford.edu/entries/qt-quantlog/> (1) implication ( $Q_1 \Rightarrow Q_2$ ),

(2) union ( $Q_3 = Q_1 \vee Q_2$ ), and

(3) intersection ( $Q_3 = Q_1 \wedge Q_2$ ).

(1) always false ( $Q_0$ ) and always true ( $Q_\infty$ )

(2) negation of question ( $?Q$ )

(3) orthogonality: if  $Q_1 \Rightarrow ?Q_2$  then  $Q_1$  and  $Q_2$  are orthogonal denoted ( $Q_1 \perp Q_2$ ).

- **quantum tetrahedron:** A quantum tetrahedron is a single tetrahedron whose geometric variables are considered as dynamical and represented by operators on a quantum state space  $\mathcal{H}$ .

- **quintessential dark matter:**

“Reinterpreting quintessential dark energy through averaged inhomogeneous cosmologies”, [astro-ph/0609315].

“...Einstein General Relativity: the cosmic quintessence emerges in the process of interpreting the real Universe in a homogeneous context.”

“...It consists in defining cosmologies that are homogeneous on large scales without supposing any local symmetry, thanks to a spatial averaging procedure. It results in equations for a volume scale factor that not only include an averaged matter source term, but also additional terms that can be interpreted as the effects of the coarse-grained inhomogeneities on the large scales dynamics. These additional terms are commonly named backreaction.”

- **RAQ:** Refined algebraic quantization.

- **radar time:** Popularized by Bondi, *radar time* is used to define hypersurfaces of simultaneity in terms of physical clocks and light pulses, as such it should be independent of the choice of coordinates.

Consider an observer travelling on path  $\gamma : x^a = x^a_{(\gamma)}(\tau)$  with proper time  $\tau$ . Define

$\tau^+(x) :=$  (earliest possible) proper time at which light ray (null geodesic) leaving the point  $x$  could intercept  $\gamma$ .

$\tau^-(x) :=$  (latest possible) proper time at which a light ray could leave  $\gamma$ , and still reach the point  $x$ .

$\tau(x) := \frac{1}{2}(\tau^+(x) + \tau^-(x))$  is the *radar time*.

$\tau(x) := \frac{1}{2}(\tau^+(x) - \tau^-(x))$  is the *radar distance*.

$\Sigma_{\tau_0} := \{x : \tau(x) = \tau_0\}$  is the observer's *hypersurface of simultaneity at time  $\tau_0$* .




Figure A.36: radartimeF. Schematic of the definition of radar time  $\tau(x)$ .

- **Raychauduri equation:** Equation that governs the expansion of an infinitesimal bundle of geodesics. There are separate versions for time-like and null geodesics.

Play a central role in the proof of the singularity theorems.

- **real formalism:** With the introduction of a generalized Wick transform to map the constraint equations of Riemannian general relativity to those of the Lorentzian theory.

- **recombination:** This is the moment, in the history of the universe, when free electrons recombine with protons to form hydrogen atoms. Matter ceases to be ionized, and decouples from electromagnetic radiation. This defines the period of decoupling, a million years or so after the Big Bang.

- **reconstruction problem:** how do we extract local geometry and dynamics from the global nature of observables. Observables are to be invariant under active diffeomorphisms. local from non-local observables.

- **reconstruction of quantum mechanics:** See quantum logic reconstruction.

- **reduced phase space:**

- **reduced phase space quantization:** Solve the constraints at the classical level. Quantize without any constraints. Given two 3-metrics one would need to know if one is a time-evolved version of the other, this requires solving for the dynamics. A reduced

phase space quantization of gravity would require finding the general solution of the classical field equations.

In the reduced phase space quantization one first performs a gauge fixing and then quantizes. That is one has to represent the algebra of gauge fixed functions equipped with the Dirac bracket as (self-adjoint) operators on a Hilbert space.

*“In her recent work, Dittrich generalized Rovelli’s idea of partial observables to construct Dirac observables for constrained systems to the general case of an arbitrary first class constraint algebra with structure functions rather than structure constants. Here we use this framework and propose how to implement explicitly a reduced phase space quantization of a given system, at least in principle, without the need to compute the gauge equivalence classes.”*

[301] based on the framework constructed in [300]

the ultimate Hilbert space must carry a representation of the Poisson algebra of Dirac observables.

• **refined algebraic quantization (RAQ):**

RAQ, as it so far formulated, cannot rigorously deal with constraints which involve non-trivial structure functions since the group averaging really requires an honest (Lie) group structure.

RAQ has actually two implementations: A heuristic version, called group averaging, and a rigorous version, using Rigged Hilbert Spaces.

• **reflection positivity:** Let  $\theta$  denote the action of the time reflection, i.e. reflection with respect to the hyperplane

$$(x^0 = 0, x^1, \dots, x^d), \tag{A.134}$$

then  $\theta g$  is the time reflection of the function  $g$ . Consider functions of the form

$$f(\phi) = \sum_{j=1}^N c_j e^{i\phi(e_j^+)}, \tag{A.135}$$

where  $c_j$  is a complex number and  $e_j^+$  are arbitrary Schwartz functions that are non-zero in the  $x^0 > 0$  half-space. Then reflection positivity of the measure  $\mu(\phi)$ , is the condition that

$$\langle \theta f, f \rangle_{L_2} = \int_{S'} (\theta f(\phi))^\dagger f(\phi) d\mu(\phi) \geq 0. \tag{A.136}$$

It is the axiom of Osterwalder-Schrader theorem that allows one to construct a physical Hilbert space of the canonical quantum field theory from the Euclidean one with a non-negative Hamiltonian acting on it.

- **Regge manifold:** A  $d$ -dimensional Regge manifold is a metric space composed by glued flat  $d$ -simplices, with matching geometry at the intersections.
- **Regge model:** One approximates a curved spacetime by a large collection of simplices glued together. The interior of every simplex is flat, the spacetime curvature resides in how the simplices are glued to each other.




Figure A.37: ReggeGlos. Gluing flat simplices to get a surface with curvature.

### Regge Lagrangian:




Figure A.38: Regge Lagrangian.

denote the lengths of its edges by  $l_i$  ( $I = 1, 2, \dots, 6$ ) (Fig E.7). The Regge action for the tetrahedron is

$$S_{Regge} = \sum_{I=1}^6 l_i \theta_i, \quad (\text{A.137})$$

where  $\theta_i$  is the angle between the outward normals of two faces sharing the  $I$ -th edge.

- **reheating:** Process whereby the period of inflationary expansion gives way to the standard hot big bang scenerio.
- **relational notion of space:** Aristotle to Decartes picture of space where described by the conjunuity of objects: there is no notion of space without matter.

• **relational interpretation of quantum mechanics:** Rovelli's relational quantum mechanics. resolves the paradox of superpositions by making it a consequence of one's point of view.

It does not try to append a reasonable interpretation to the quantum mechanics formalism, nor to postulate as yet unknown new physics, rather it attempts to uncover the physical meaning quantum mechanics. One drops the assumption that quantum states are observer independent.

In the relational interpretation, the very notion of state of a physical system should be considered meaningless unless it is understood to be relative to another system that plays the role of one possible observer. No implication that the observer is human or has any other a priori special properties only that it has the possibility of interacting with the 'observed' system.

It is the specification of such an observer that allows the ascription of a state to a system to make sense. So the description the theory provides is complete because it exhausts everything that can be said about the world.

A physical quantity has value with respect to certain observer.

The cat finds himself in one or other of the possible outcomes while at the very same time according to the person outside the box the cat is in a quantum superposition. People say the two can't be true at the same time so we have a paradox. However this discord does not impinge on you because when to different observers can ask the same question they get the same answer.

That the photon has been absorbed (or not) according to the cat isn't to say that it has been (or has not been) absorbed as far as anyone else is concerned - it is not observer independent, that is it is not an absolute property of the system. In this interpretation there is no need for special systems that collapse the wavefunction for everyone else's benefit.

There is a long history in physics of progress being made by abandoning the belief that there are preferred systems and by 'relationalizing' notions that were previously thought to be absolute. The description provided by the theory exhausts everything that can be said about the world.

What is stopping us from taking this as a description of reality is a philosophical prejudice and nothing more. We should adjust our philosophical prejudices to what nature is telling us. We shouldn't be trying to fit reality into our preconceived notions about the world.

<http://plato.stanford.edu/entries/qm-relational/>

consistent discretizations

context created by the presence of an observer

different descriptions as there are as there are contexts, but with interrelations between them so that if two observers can ask the same question they get the same answer.

- **Relational quantization *a la* Page-Wootters:** [47]

- **relational quantum cosmology:**

[29] Slogan: “Many quantum states to describe one universe, not one state describing many universes”.

- **relativistic spin networks:**

- **renormalization:**

renormalized mass and charge values are inserted into the theory as unexplained parameters. Coupling constants of various kinds and masses of the basic quarks and leptons, the Higgs particle, etc. need to be specified.

Any acceleration of a charged particle entails the acceleration of part of the charged particles surrounding it, so the mass  $m$  that you observe is the mass  $m_0$  of the particle plus a correction  $\Delta m$ , due to its interaction with the cloud of charged particles. Procedure of replacing infinities with experimentally determined values. a quantum field theory whose Lagrangian has a few free parameters - masses and charges and so. These quantities are referred to as the “coupling constants”.

with obvious physical meaning cannot be calculated without incurring divergencies.

Renormalization occurs in evaluating physical observable quantities which in simple terms can be written as formal functional integrals of the form

$$\int [\mathcal{D}\phi] e^{\mathcal{L}(\phi, \partial\phi)} A(\phi, \partial\phi) \tag{A.138}$$

where  $\mathcal{L}$  is composed by two parts, a part quadratic in  $\phi$  and  $\partial\phi$  term,  $\mathcal{L}_0$ , and an ‘interaction’ term,  $\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_{int}$ . Computing such an integral in perturbative terms leads to a formal power series, each term  $G$  of which is obtained by integrating a polynomial under a Gaussian  $e^{\mathcal{L}_0}$ :

$$\int [\mathcal{D}\phi] e^{\mathcal{L}_0(\phi, \partial\phi)} (1 + g\mathcal{L}_{int}(\phi, \partial\phi) + \frac{g^2}{2!}[\mathcal{L}(\phi, \partial\phi)]^2 + \dots) A(\phi, \partial\phi) \tag{A.139}$$

Such Feynmann diagrams  $G$  are given by multiple integrals over spacetime or, upon Fourier transformation, over momentum space, and are typically divergent in either case. The renormalization technique consists in adding counterterms  $\mathcal{L}_G$  to the original Lagrangian  $\mathcal{L}$ , one for each diagram  $G$ , whose role is to cancel the undesired divergencies.

In the case of a renormalizable theory, all the necessary counterterms can be obtained by modifying the numerical parameters of the original Lagrangian.

considered to be ad-hoc

Kreimer there is a Hopf algebra underlies quantum field theory renormalization, and has led to a new approach to perturbative renormalization under development.

The antipode of a diagram is related to the counterterms necessary to subtract its divergencies.

**renormalization - background independent:** The regularized Hamiltonian operator is independent of the regularization parameter  $\epsilon$  to first order - the regularization parameter has disappeared from the expression! Hence no renormalization is involved! The reason why the regularization parameter does not appear is, roughly speaking, because the shrinking process,  $\epsilon \rightarrow 0$ , can be compensated for by an active spacial diffeomorphism.

Tentative ideas have been formulated by Markopoulou [122], [123] and Oeckl [124].

**renormalization a la Fontini-Markopoulou** [122], [123]

$$\gamma_1 \subset \gamma_2 \quad \gamma_2 \subset \gamma_1 \quad \gamma_1 \cap \gamma_2 = \emptyset$$

let  $\gamma$  denote a proper sublattice of  $\Gamma$ , namely  $\gamma \neq e$  and  $\gamma \neq \Gamma$ . We call the lattice that is left after we “cut out”  $\gamma$  the **remainder** and denote it  $\Gamma/\gamma$ .

$$\Delta(\gamma_p) = \gamma_p \otimes e + e \otimes \gamma_p \tag{A.140}$$

These are the **primitive elements** of the Hopf algebra.

The **counit** is an operation which annihilates every lattice except  $e$ .




Figure A.39:

**renormalization a la R. Oeckl** [124]

- **renormalization group:** A method designed to describe how the dynamics of a system change as we change the scale at which we probe it.

- **renormalized charge:** bare charges divergent, absorption of infinities into a redefinition of the charges. screening effects (see effective charges).

- **Ricci identity:**

$$2\nabla_{[a}\nabla_{b]}V_c = R_{abcd}V^d \quad (\text{A.141})$$

- **rigged Hilbert space - also called the Gel'fand triple:** In infinite dimensional Hilbert space there can be operators that have a continuous spectrum, (the most simple example would be that of a free electron which has a continuous momentum spectrum). The corresponding eigenvectors are not normalizable, and hence cannot lie inside the Hilbert space  $\mathcal{H}$ , (the eigenvector of the momentum operator of a free electron is a plane wave  $e^{ikx}$ , which isn't normalizable). The rigged Hilbert space approach to dealing with such operators on or a Gel'fand triple  $\{\Phi, \mathcal{H}, \Phi'\}$ , in which the Hilbert space  $\mathcal{H}$  appears as a linear subspace  $\Phi'$  which is not itself a Hilbert space. Eigenvectors associated with a continuous spectrum belong to  $\Phi'$  but are no members of its subspace  $\mathcal{H}$ . The space  $\Phi'$  is constructed as a type of dual of another vector space  $\Phi$  which is a subspace of  $\mathcal{H}$ . Thus we have  $\Phi \subset \mathcal{H} \subset \Phi'$ .

- **Riemannian geometry:** A generalized geometry that has the property of being locally flat; that is, in a sufficiently small region, a Riemann geometry can be approximated by a Euclidean or Minkowski geometry.

- **Robertson-Walker metric:** The metric that describes an isotropic and homogeneous cosmological space time.

- **Rovelli, Carlo:**



Figure A.40: Carlo Rovelli.

- **Sandwich Conjecture:** The conjecture that, given two spatial metrics  $q$  and  $q'$  on two hypersurfaces then there is unique metric that will interpolate between them, up to

gauge.

- **scale factor:** The quantity that describes how length change in the expanding (or contracting) universe.

- **scattering amplitudes:**

There is no given *a priori* geometry here, as is the case in background dependent theories. The geometry is encoded in the quantum state itself, not given a priori as in conventional QFTs.

$$\int \mathcal{D}\varphi_I \int \mathcal{D}\varphi \tag{A.142}$$

how can any physics possibly come out of the setup here if the amplitude doesn't change when deforming the boundary surface - that is, until you realize that the metric is one of the boundary fields!

$$\langle \Psi_1 | \Psi_2 \rangle = \int \mathcal{D}\varphi_m \Psi_1^*[\varphi_m] \Psi_2[\varphi_m], \quad |\Psi_1 \rangle, |\Psi_2 \rangle \in \mathcal{H}_{\Sigma_m}. \tag{A.143}$$

The physical information is contained in the spectrum of the boundary operators of partial observables.

- **scattering matrix:** The scattering matrix allows the to calculate experimentally observable scattering cross sections.

- **Schödinger equation:** The equation that describes the evolution of a nonrelativistic wavefunction.

- **Schödinger picture:**

- **Schwarzschild radius:** The radius of the event horizon of a nonrotating black hole of mass  $M$ , equal to  $2GM/c^2$ .

- **secondary constraint:** Dirac bracket substituting for the Poison bracket

- **second-quantization:** The passage from a one particle system with an arbitrary number of particles subject to the Bose/Fermi statistics is called second-quantization.

$\hat{\Psi}(\mathbf{x})$  does not represent the actual quantum state it is the operator that creates a new particle having the one-particle wavefunction  $\psi(\mathbf{x})$ , by application on the ground state  $|0 \rangle$  (the state representing the absence of any particles).

- **self organized criticality:** Self organized critical systems are statistical systems that naturally evolve without fine tuning to critical states in which correlation functions are scale invariant. The earliest example of such a system is the sandpile of Bak, Tang and

Wiesenfeld [258]. Since then, many such systems have been studied, including models of phenomena as diverse as biological evolution, earthquakes, astrophysical phenomena and economics [259].

It has been suggested that the emergence of classical spacetime from a discrete microscopic dynamics may be a self-organized critical process, [257].

- **semi-classical quantum gravity:** Semi-classical quantum gravity refers to the setting in which a classical background geometry is coupled to the quantum matter field through the semiclassical Einstein equations:

$$G_{\mu\nu} = -8\pi \langle T_{\mu\nu} \rangle .$$

It is a perturbative approach where there is a well-defined classical background geometry about which the quantum fluctuations are occurring.

- **separating:** Functions of phase space  $\mathbb{R}^{2n}$  - observables. the energy and momentum functions on phase space is sufficient know to find out a which particular point in phase space the system is at.

these functionals form a separating set on  $\mathcal{A}/\mathcal{G}$ : if all the  $T_\alpha$  assume the same values at two connections, they are necessarily gauge related.

- **shadow states:** It would help in assesing the viability of LQG providing the correct semi-classical limit if we closed the gap between its one dimension excitation description with the usual Fock space description, in the low energy limit cutoff.

The problem with the polymer representation is that,  $Cly^*$ , the space containing the solutions to the quantum constraint equations, does not have a physically justified inner product; a definite Hilbert space structure is not yet available. While the kinematic states belong to  $Cyl$ , the physical states belong to  $Cly^*$ . Therefore, semi-classical states, capturing the low energy physics, should also be in  $Cly^*$ . Can we still test a candidate state in  $Cyl^*$  for semi-classicality at this stage without acces to expectation values? One uses the notion of *shadow states* to do semiclassical analysis in the polymer representation.

Shadow states use graphs as ‘probes’ to extract physical information from elements  $(\Psi|$  of  $Cly^*$ . More precisely, one ‘projects’ each  $(\Psi|$  to an element  $|\Psi_\gamma^{shad} \rangle$  in  $Cyl_\gamma$ , a so-called shadow state, and analyses properties of  $(\Psi|$  in terms of its shadows  $|\Psi_\gamma^{shad} \rangle$ . Each shadow captures only a part of the information contained in our state, but the collection of shadows can be used to determine the properties of the full state in  $Cly^*$ .

- **simplicial spin-foam:** Standard spin-foam models as formulated up to now are piecewise flat geometries defined on piecewise linear manifolds. For LQG canonical theory to match these models it would have to be restricted to the category of piecewise linear manifolds and piecewise linear spin-networks. A suitable generalisation of simplicial spin-foam models to lift this restriction is linear 2-cell spin-foam models.

- **simplicity constraints:** Imposing the simplicity constraints turns the topological BF theory into (non-topological) 4-dimensional gravity.
- **simple regions:** A simple region of spacetime  $\mathcal{M}$  is a geodesically convex set with compact closure, whose boundary is diffeomorphic to the three sphere  $S^3$ .
- **singularity:** the inevitable presence of singularities in general relativity of the classical gravitational being always attractive.

A singularity is a point in a spacetime which the curvature becomes infinite. Robertson-Walker, Schwarzschild, Reissner-Nordstrom and Kerr are examples of solutions contain points of infinite curvature. The presence of infinite curvature lets itself know by timelike or null geodesics coming to a full stop. In the context of the singularity theorems, the working definition of a singularity is a kind of incompleteness of the space-time under consideration, more precisely, it is an obstruction of some sort to timelike or null geodesics from being indefinitely extendible. Investigate the nature of singularities.

We require a smooth Lorentz metric. You cannot have point on the manifold with singular metric. So the points where this can happen are excised.

- **singularity theorems:** Proved by Hawking and Penrose. The singularity theorems are based on very powerful indirect arguments which show that black hole and cosmological singularities are generic in classical general relativity. Gives us confidence in the big bang. The classical theory predicts its own breakdown. One of the main motivations for attempting to quantize general relativity.

A feature of the singularity theorems is that they do not directly show the existence of black holes. Instead they show that spacetime is geodesically incomplete so that the worldline of an observer comes to an end and cannot be extended. The obstruction to extending the worldline is some kind of singularity, but this might be rather mild and need not correspond to a black hole.

If there is generically in spacetime a closed, trapped region, energy certain positivity and if gravity remains attractive a singularity is inevitable. the conclusion is that there is to be some some sort of obstruction to timelike or null geodesics from being indefinitely extendible.

- **slow-roll approximation:**

from gr-qc/0511007:

The mechanism, proposed in [], considers so called slow-roll approximation, when the inflation rolls down its potential hill towards the potential minimum, but, at the same time, remaining far away from the minimum. It is exactly in the vicinity of the minimum of the potential, where the inflation stops.

- **Smarr's formula:** Smarr's formula translates black hole mass  $M$  to its angular momentum  $J$ , angular velocity  $\Omega_H$ , surface gravity  $\kappa$ , and surface area  $A$ :

$$M = 2\Omega_H J + \frac{\kappa A}{4\pi}. \quad (\text{A.144})$$

- **Smolin, Lee:**



Figure A.41: Lee Smolin.

- **Smolin’s Darwinian hypothesis:** Smolin put forward the proposal that the universe fine tunes the values of its physical constants through a Darwinian selection process [184]. Every time a black hole forms, a new universe is developed inside it that has different values for its physical constants from the one in its progenitor.

[125] not just based on “*authors intuition*”

- **space of solutions:** The space of solutions are taken to be the phase space or space of states. This definition is independent of any special time choice so that it is manifestly covariant.

- **spacial averaging problem:**

- **spacial diffeomorphism invariance:**

- **spacetime singularity** The working definition of a spacetime singularity is a point in spacetime beyond which null or timelike geodesics cannot be extended. We mean the singularity

- **special relativity:**

- (i) **The principle of relativity.** The laws of physics are the same for all inertial reference frames.

(ii) **The constancy of the speed of light.** The speed of light in a vacuum is the same for all inertial observers irrespective of the motion of the source.

• **Speziale, Simone:** Work on spin foams and the graviton propagator in loop quantum gravity.



Figure A.42: Simone Speziale.

• **spin foam:** Spin foam models are an attempt to describe the geometry of spacetime at the quantum level. They give a construction of transition amplitudes between initial and final spacial geometries labelled by spin networks.

???the spin foam is independent of any special time choice so that it is manifestly covariant???

Spin foams can be understood to be dual to triangularizations.

In this sense the physical state is associated with an entire history of the system, rather than any point on that history.

• **spin network:** A spin network is a generalization of a knot or link. It is a graph embedded in space, with edges carrying labells corresponding to an irreducible representation of a Lie group, and vertices with labells corresponinding to interwining operators.

• **spin network state:** They diagonalize geometric operators and describes quantized three geometry. They provide a complete orfonormal basis for the kinematic Hilbert space. Using spin networks we are working with finite dimensional spaces that have the structure of Hilbert space of spin systems - the same mathematics at work as in ordinary quantum mechanics of angular momentum but with completely different interpretation. They are natural candidates for cylindrical functions in the construction of well defined measures of integration. The scalar product has a very simple form  $\langle \Phi_\alpha \Phi_\beta \rangle = \delta_{\alpha\beta}$ . This make easy at the time of calculations.

Spin networks states provide a natural decomposition of gauge-independent kinematic Hilbert space,  $\tilde{\mathcal{H}}^0$ , into finite dimensional subspaces each of which can be identified with the space of states of a spin-system. This simplifies various constructions and calculations enormously.

Using spin networks we are working with finite dimensional spaces that have the structure of Hilbert spaces of spin systems.

- **spinor:** It is well known that the Pauli matrices satisfy the commutation relations of the rotation algebra,

$$[\tau_i, \tau_j] = \epsilon_{ijk} \tau_k \tag{A.145}$$

They therefore are a representation of the rotation algebra; they generate infinitesimal rotations on a two component vector - this is the 2-component spinor. Finite rotations are obtained by the exponentiating of (E.7),

$$\begin{aligned} \hat{U}_R &= \exp(-i\phi \cdot \tau) = \exp(-i\frac{1}{2}\phi \mathbf{n} \cdot \tau) \\ &= \mathbf{I} \cos(\frac{1}{2}\phi) - i\mathbf{n} \cdot \tau \sin(\frac{1}{2}\phi). \end{aligned} \tag{A.146}$$

The defining property of a spinor is the way it transforms under rotations (or Lorentz transformations depending on the context).

- **Standard model:** particle physics.  $SU(3) \times SU(2) \times U(1)_Y$ . The Standard Model is not a mathematically consistent theory. It has had undeniable success perturbative calculations of collision cross section, the numerical analysis of particle spectra etc.

- **State:**

There is a second use of this term. A collection ‘observables’ and is capable of being in certain ‘states’. We can define the state of a system as knowledge of the expected values of the observables, that is, a state is an assignment of an expected value of each observable.

A second use of this term is *an expectation value functional*. This is to be chosen at some point in the quantization.

$$F_{\text{Sch}}(W(\zeta)) := e^{-\frac{1}{2}|\zeta|^2}. \tag{A.147}$$

- **state sum model:** [411]

- **Stone-von Neumann uniqueness theorem:**

$$\begin{aligned}\{q, p\} &= 1, \\ \{q, q\} &= \{p, p\} = 0.\end{aligned}\tag{A.148}$$

$$[q, p] = i\hbar, \quad [q, q] = [p, p] = 0.\tag{A.149}$$

Now we find concrete representation of this algebraic relation. Schrödinger's wavefunction representation

$$\hat{q}\psi(q) = q\psi(q), \quad \hat{p}\psi(q) = -i\hbar\frac{d\psi(q)}{dq}\tag{A.150}$$

$$[\hat{q}, \hat{p}]\psi(q) = (-i\hbar q\frac{d}{dq} + i\hbar\frac{d}{dq}q)\psi(q) = i\hbar\psi(q)\tag{A.151}$$

Heisenberg's matrix mechanics every observable physical quantity, that is energy, position, momentum, angular momentum, is described by an operator represented as a matrix.

Schrödinger's wavefunction representation and Heisenberg's matrix mechanics describe the same physics. Are essentially unitary equivalent.

In quantum field theory there is no analog of the Stone-von Neumann uniqueness theorem. Hence, there are an infinite number of inequivalent representations of the Poisson bracket algebra. These inequivalent representations can be thought of as different "phases", which have different physics.

The Fock representation the natural generalization of the Schrödinger representation to infinitely many degrees of freedom. Quantum field theories written in the Fock representation by theoretical physicists.

- **string theory:** all known particles and their interactions are supposed to emerge as certain modes of excitation of and interactions between quantized strings.

String theory attempts to produce a theory of everything, including quantum gravity that will have general relativity as part of the classical limit.

a little "far-out". String theory is a perturbative approach depending on the choice of background metric. background independent M-theory for strings, Witten using twistor space methods in which physics happens in twistor space and spacetime is a secondary construction.string field theory.

*We find that the massless closed quantum states include one-particle graviton states, making string theory a quantum theory of gravity!*

Except string theory can only be defined over stationary spacetimes, which have measure zero in the space of solutions to Einstein's equations - a timelike Killing vector field is needed to have spacetime supersymmetry, without it you get unphysical tachions in the spectrum. dynamical theory over spacetime confused with a dynamical theory of spacetime.

A necessary condition for a perturbative theory to be consistent is that the two dimensional world sheet quantum field theory that defines the theory be conformally invariant. This means that the conformal anomaly on the two dimensional worldsheet vanishes. To leading order in  $l_{string}$  this condition is equivalent to the Einstein equations of the background manifold.

The so called Pohlmayer string can be quantized in any number of dimensions, including four dimensional Minkowski space [?].

Background dependency: (arXiv:hep-th/9808192)

in which the definitions of the states, operators and inner product of the theory require the specification of a classical metric geometry. The quantum theory then describes quanta moving on this background. The theory may allow the description of quanta fluctuating around a large class of backgrounds, but nevertheless, some classical background must be specified before any physical situation can be described or any calculation can be done. All weak coupling perturbative approaches are background dependent, as are a number of non-perturbative developments. In particular, up to this point, all successful formulations of string theory are background dependent.

String theory only studies semiclassical effects in fixed classical spacetime backgrounds.

- **strongly causal:** There are several types of causality violations that can be considered, other than the existence of closed timelike curves. Strong causality is an important example. Roughly speaking, strong causality is said to fail if an arbitrarily small perturbation of the metric will result closed timelike curves, that is, the chronology condition is almost violated. Strong causality plays a key role in the Hawking-Penrose singularity theorem. Important there is the result: Let  $\mathcal{M}$  be a spacetime satisfying: (1) There are no closed trips. (2) Every endless null geodesic in  $\mathcal{M}$  contains a pair of conjugate points. Then  $\mathcal{M}$  is strongly causal.

see future-distinguishing spacetimes, stably causal

- **supergravity:**

Bosonic and fermionic Feynman diagrams cancel each other out.

- **superselection:** A representation is irreducible if the Hilbert space does not split into orthogonal subspaces preserved by the action of  $\mathcal{A}$ . Equivalently, there must exist in the Hilbert space a cyclic vector whose image under the action of  $\mathcal{A}$  is dense in the Hilbert space. Physically this requirement rules out the existence of superselection rules: superselection sectors can be identified with the irreducible sectors of a reducible representation.

The representation should be irreducible on physical grounds (otherwise we have superselection sectors implying that the physically relevant information is already contained in a closed subspace).

- **superspace:** the collection of all possible three-metrics (nothing to do with supersymmetry)

- **supersymmetry:**

Supersymmetry requires fermions and bosons to come in pairs consisting of one of each, with the same mass. This is not observed in nature.

- **surface gravity:** **Surface gravity** is the acceleration of a static particle near the horizon as measured at spacial infinity.

$$\kappa = -\frac{1}{2}(\nabla^a \xi^b)(\nabla_a \xi_b)|_{\mathcal{N}} \quad (\text{A.152})$$

- **symmetry:** In quantum mechanics group transformations that leaves invariant key structures in the theory - Hilbert space, algebra of observables, etc.

An unitary operator on Hilbert space preserves all scalar products, i.e.,

$$\langle \hat{U}\psi, \hat{U}\phi \rangle = \langle \psi, \phi \rangle \quad (\text{A.153})$$

- **symplectic form:**  $\omega_{\mu\nu}$  is antisymmetric and also has the properties

$$\omega^{\mu\nu} = \{\xi^\mu, \xi^\nu\} \quad (\text{A.154})$$

(i) It is antisymmetric  $\omega_{\mu\nu} = -\omega_{\nu\mu}$ ,

(ii) is non-degenerate, i.e.,  $\omega_{\mu\nu}v^\nu = 0$  implies  $v^\mu = 0$ , that means that the inverse matrix  $\omega^{\mu\nu}$  exists,

(iii) it is closed, i.e.,  $\partial_\mu \omega_{\nu\gamma} + \partial_\nu \omega_{\gamma\mu} + \partial_\gamma \omega_{\mu\nu} = 0$  which is equivalent to the fact that the Poisson bracket satisfies the Jacobi identity.

$$\omega = \frac{1}{2}\omega_{\mu\nu}d\xi^\mu \wedge d\xi^\nu \quad (\text{A.155})$$

- **tempered distributions:** they are continuous (or equivalently - bounded) linear maps from the Schwarz space of functions to complex numbers. In other words, it is the topological dual space to the of  $\mathcal{S}(\mathbf{R}^n)$

- **tensors:** The point about tensors is that when we want to make statement we do not wish it to hold in just one coordinate system but rather all coordinate systems.

- **tensor calculus:**

- **test particle:** is an idealized point particle with energy and momentum so small that its effects on spacetime are negligible.

- **thermal clocks:** Clocks that change linearly with respect to thermal time. (Rovelli)

- **thermal field theories:**

- **thermal time hypothesis:** The dynamical laws of generally covariant systems determine correlations observables, but they don't single out any of the observables as that we would call "time".

two equivalent ways of describing the time flow: either the flow in state space (Schrödinger picture), or (generalized Heisenberg picture)

So we say time flow is as a one-parameter group of automorphisms of the algebra of observables.

the physical basis of time [290] [291].

$$\mathcal{H} = -kT \ln D \tag{A.156}$$

The thermal states of a field theory are characterized by have a correlation function, the KMS condition (A.89)

- **thermodynamics:** Thermodynamics deals with large systems in terms of macroscopic observables alone.

know as the zeroth, first and second laws

**zeroth:**

**first:**

**second:**

Bekenstein suggested that the second law of thermodynamics should be extended in the presence of black holes.

- **Thiemann, Thomas:**

- **Thomson scattering:** When a photon scatters off a free electron, in the nonrelativistic case  $\hbar f \ll m_e c^2$ .

- **tidal force:**



Figure A.43: Thomas Thiemann.

- **top-down:** the is required. Loop quantum gravity is an example. One of the biggest challenges for the theory, most be established before it can be claimed to be a viable theory of quantum gravity.

- **time:** ??

- **timelike membranes:** Their definition is the same as a dynamical horizons except that they are timelike instead of spacelike. [149]

- **topological quantum field theories:**

They have a finite number of degrees of freedom so are really mechanical systems disguised as field theories. they posses certain features one expects of a quantum theory of gravity.

- **trace class operator:** nuclear operator.

- **trapped surfaces:** A trapped surface  $\theta_{(\ell)} < 0$  and  $\theta_{(n)} < 0$

**marginally trapped surface:** A marginally trapped surface  $\theta_{(\ell)}, \theta_{(n)} \leq 0$

An **outer marginally trapped surface**  $\theta_{(\ell)} \leq 0$ . is the boundary of a three-dimension volume whose expansion of the outgoing family of null geodesics orthogonal to  $S$  is everywhere non-positive,  $\theta_{(\ell)} \leq 0$ .

An apparent horizon

A closed, spacelike, two-surface is an apparent horizon if it is the outermost marginally trapped surface.

**trapped boundary a la Hayward:**

**trapped surface a la Penrose:**

**closed trapped surface:**

- **twisted geometry:** A basis of states for LQG is given by the spin network states. A spin network state has support on a graph  $\Gamma$  and determines a 3d “quantum geometry”; they are eigenvectors of geometric operators which determine the intrinsic geometry. The extrinsic geometry (analogous to the conjugate momentum) is completely “spread out”, due to the Heisenberg uncertainty principle. To make contact with a semiclassical description of space, we can consider coherent states peaked (but not sharp) on both the intrinsic and extrinsic geometry. A twisted geometry is a specific choice of “interpolating geometry”, chosen among discontinuous metrics. To any graph and any holonomy-flux configuration, we can associate a twisted geometry: a discrete discontinuous geometry on a cellular decomposition space into polyhedra. Thanks to this result, the phase space of LQG on a graph can be visualised not only in terms of holonomies and fluxes, but also in terms of a simple geometric picture of adjacent flat polyhedra.

- **twistor theory:** invented by Sir Roger Penrose. : entire light-rays are represented as points, and events by entire Riemann spheres. Twistors defined in terms of a pair of spinors. Twistors are the coordinates of twistor space. In twistor theory spacetime is a secondary concept. Witten is keen on twistor theory in developing background-independent string (M-)theory. Witten found that a lot of the beautiful results of twistor theory did not extend from four dimensions to higher dimensions. If Thiemann’s LQG-string theory stands up to the critical examination, in four dimensions there could be possibility of applying the powerful results of twistor theory to a background independent string theory?? [16]. Seminar given by Penrose (2004) *Goals and Achievements of Twistor Theory*, [444].

- **2+1 quantum gravity:** Witten Chern-Simons. See ??.

In a vacuum in three-dimensions the vanishing of the Ricci tensor implies the Riemann tensor vanishes

$$R_{ab} = 0 \implies R^a{}_{bcd} = 0 \tag{A.157}$$

so the solution of the equations of motion is that curvature vanishes.

This condition, which is imposed *a priori* in Einstein’s formulation of general relativity, is seen to be part of the equations of motion.

$$\mathbf{R}^3 \otimes_S SO(3). \tag{A.158}$$

- **uncertainty relations:**

$$\Delta A \Delta B \geq \frac{1}{2} | \langle [\hat{A}, \hat{B}] \rangle |, \quad (\text{A.159})$$

where  $\Delta A = \sqrt{\langle \hat{A}^2 \rangle - \langle \hat{A} \rangle^2}$ .

- **unification:** String theory.

Could degrees of freedom corresponding to matter arise in some natural way from those of geometry?

Chakraborty-Peldan unified model [338]

*“...If this difficulty could be overcome and the model could be made to include Fermions via supersymmetry it would become a viable, elegant and mathematically well defined way of having a unified theory of quantum fundamental interactions.”*

*Quantum Gravity and the Standard Model*

Discrete quantum gravity: a mechanism for selecting the value of fundamental constants [328].

- **uniform discretization:** A particularly promising version of “consistent discretizations”.

Several approaches to the dynamics of loop quantum gravity involve discretizing the equations of motion. Discretized theories are problematic since the first class algebra of constraints of the continuum theory become second class upon discretization. If one treats the second class constraints normally, the resulting theories have very different dynamics and number of degrees of freedom than those of the continuum theory. It is therefore questionable how these theories could be considered an appropriate starting point for quantization and the definition of a continuum theory through a continuum limit. The uniform discretization approach is a proposal for the quantization of constrained systems which could overcome these difficulties and construct the correct quantum continuum limit.

- **uniqueness theorems:** See black hole uniqueness theorems.
- **unitarity:** to preserve probabilities

Unitarity implies that expectation values of gauge invariant observables does not depend on the gauge or frame of reference.

- **universe:** That which contains and subsumes all the laws of nature, and everything subject to those laws; the sum of all things that exists physically, including matter, energy, physical laws, spacetime.

- **Unruh effect:** An observer who accelerates uniformly through flat empty space will observe a thermal bath of particles, at a temperature given by their acceleration. This

means that a state which is empty according to one observer will not be empty according to an accelerating observer, and hence demonstrates that the concept of “vacuum” must be observer dependent. Discovered by Unruh [385] and Davies [386].

- **vacuum:**

$$|0 \rangle \tag{A.160}$$

define the vacuum state as follows

$$a(k)|0 \rangle = 0 \tag{A.161}$$

where

$$W(q_1, q_2, \dots, q_n) = \langle q_1, q_2, \dots, q_n | 0 \rangle \tag{A.162}$$

where  $\langle q_1, q_2, \dots, q_n |$  are eigenstates of observable quantities.

$$\lim_{t \rightarrow \infty} W(\alpha, -it, \alpha', 0) \rightarrow H_0(\alpha) e^{-E_0 t} \overline{H_0(\alpha')} \tag{A.163}$$

$$\lim_{t \rightarrow \infty} e^{E_0 t} |0_{-it} \rangle = |0_M \rangle \langle 0_M | \tag{A.164}$$

- **vacuum expectations:**

$$G(x, y) = \langle 0 | a(y) a^\dagger(x) | 0 \rangle \tag{A.165}$$

- **Vaidya:** The spherically solution to Einstein’s equations with a null fluid as source.

describe the formation of a black hole through infalling null dust.

Simplest examples of spacetimes admitting dynamical horizons.

- **Varadarjan, :**

- **Varadarjan isomorphism:**

$$\sum_i a_i W_{\alpha_i}[A] = 0. \tag{A.166}$$

$$\sum_i a_i \alpha_i = 0. \tag{A.167}$$

- **volume operator:**  $\hat{\mathcal{V}}_{AL} \hat{\mathcal{V}}_{RS}$

- **von Neumann, John:** Developed rigorous mathematical foundations to quantum mechanics. Important in quantum gravity where experimental guidance, so far, is missing. There is the Stone-von Neumann uniqueness theorem - a version of which was developed for LQG to prove the uniqueness of the kinematic representation of the holonomy-flux algebra. Developed the infinite tensor product Hilbert space, applied by Thiemann et al to quantum gravity in the semiclassical analysis and AQG.




Figure A.44: John von Neumann.

[99], [100]

- **weak anthropic principle:** The weak anthropic principle is the principle whereby the existence of life explained by random selection from an ensemble of universes with differing properties.

- **weave states:** A superposition of spin networks that mimic a given classical spacetime metric. Spin network states are eigenstates of  $E$ , so they are like "momentum" eigenstates. So a weave state is maximally spread in the configuration variable  $A^a(x)$ . Hence they can not represent semiclassical states, just as  $e^{ik \cdot x}$  can not represent a semiclassical wavefunction of a particle because measuring momentum precisely results in complete uncertainty in position.

- **weak cosmic censorship:**

1.  $\mathcal{I}^+$  and  $\mathcal{I}^-$  are complete;
2.  $I^-(\mathcal{I}^+)$  is globally hyperbolic.

- **Weiner-measure:** Brownian
- **Weyl algebra:** Translation  $V(\mu)$
- **Weyl tensor:** The Weyl tensor is the conformally independent part of the curvature tensor. It is the part of the curvature tensor describes the local degrees of freedom of the gravitational field.
- **Wheeler, John:**



Figure A.45: John Wheeler.

- **Wheeler-DeWitt equation:** the Hamiltonian constraint

all the dynamics is contained in constraints.

- **Wick:**

We take spacetime (or manifold?) to Riemannian, but physically, the spacetime is Lorentzian. We pass from one to the other through a Wick rotation.

See the generalized Wick rotation.

- **Wightman axioms:** A list of axioms taken as a blue-print of the general concept of a Quantum Field Theory ([gr-qc/0512053]). Here for special relativistic field theory.

Generalised by Thiemann *et al* in [] to generally covariant theories such as Quantum gravity.

i) There exists a Hilbert space  $\mathcal{H}$  with a dense domain  $\mathcal{D} \subset \mathcal{H}$ , so that all  $\Phi(F)$  are well defined operators on  $\mathcal{D}$ , and  $\Phi(F)^* = \Phi(\bar{F})$ .

ii)  $F \mapsto \Phi(F)$  is complex linear and suitably continuous.

iii) Covariance:

There is on  $\mathcal{H}$  a unitary representation

$$U(L)\Phi(F)\Phi(L)^* = \Phi(F \circ L^{-1}) \quad (\text{A.168})$$

with  $U(L)\mathcal{D} \subset \mathcal{D}$ .

iv) Locality, or relativistic causality:

If the *supports* of the test function  $F_1$  and  $F_2$  are causally separated, the corresponding field operators commute

$$[\Phi(F_1), \Phi(F_2)] = 0. \quad (\text{A.169})$$

v) spectrum condition/positivity of the total energy:

Writing  $U(1, a) = e^{iP_\mu a^\mu}$ , it holds (for expectation values) that

$$P_0^2 - P_1^2 - P_2^2 - P_3^2 \geq 0, \quad P_0^2 \geq 0. \quad (\text{A.170})$$

vi) Existence and uniqueness of the vacuum:

There exists  $\Omega \in \mathcal{D}$ ,  $\|\Omega\| = 1$ , so that  $U(L)\Omega = \Omega$  and this vector is uniquely determined up to phase factor.

vii) Cyclicity of the vacuum:

The domain  $\mathcal{D}$  is spanned by vectors of the form

$$\Omega, \Phi(F)\Omega, \Phi(F_1)\Phi(F_2)\Omega, \dots \quad (\text{A.171})$$

• **Wigner function:** The Wigner function of a semi-classical state is peaked along the phase-space trajectory.

• **Wilson loop:** The trace of the holonomy.

• **Wilson renormalization:**

Can be used even if theory not renormalizable.

$$\int \mathcal{D}[\phi] \tag{A.172}$$

• **Winkler, Oliver**




Figure A.46: Oliver Winkler.

• **world function:** The world function  $\Omega(x_A, x_B)$  is defined as half the squared geodesic distance between two points  $x_A$  and  $x_B$ .

• **Winicour solutions:** Winicour J 1975 *J. Math. Phys.* **16** 1805

• **Witten model:**

[387] E. Witten, “(2 + 1)-dimensional gravity as an exactly solvable system”, *Nucl. Phys.* **B311**: 46, (1988).

• **Yang-Mills:** There is a \$1,000,000 mathematical prize for the consolidation of Yang-Mills and special relativity - does consolidation within a background-independent theory with quantum gravity coupled to the standard model count? Or specifically, background dependent, Minkowski spacetime Yang-Mills theory?

Smolin said

“someone might earn a Clay prize by rigorously constructing quantum Yang-Mills within LQG. It will certainly not be me, but there are people working on exactly that program. The conjecture is that background independent QFTs are more likely to exist rigorously in 3+1 dimensions than Poincare invariant QFTs.”

$$\emptyset, \circlearrowleft, \forall, \exists, \ell, \wp, \Rightarrow, \uparrow, \cong, \star, \simeq, \alpha, \subseteq, \gg, \ll, \supset, \not\subset, \iota, \jmath, \wp \tag{A.173}$$

